



## Probability Estimates for Multiparameter Brownian Processes

Richard F. Bass

*The Annals of Probability*, Vol. 16, No. 1. (Jan., 1988), pp. 251-264.

Stable URL:

<http://links.jstor.org/sici?sici=0091-1798%28198801%2916%3A1%3C251%3APEFMBP%3E2.0.CO%3B2-H>

*The Annals of Probability* is currently published by Institute of Mathematical Statistics.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ims.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is an independent not-for-profit organization dedicated to and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## PROBABILITY ESTIMATES FOR MULTIPARAMETER BROWNIAN PROCESSES<sup>1</sup>

BY RICHARD F. BASS

*University of Washington*

Let  $F$  be a distribution function on  $[0, 1]^d$ , and let  $W_F$  be the Gaussian process that is the weak limit of the empirical process determined by  $F$ . If  $G$  is a function on  $[0, 1]^d$ , upper and lower bounds are found for  $P(\sup_{t \in [0, 1]^d} |W_F(t) - G(t)| \leq \epsilon)$ .

**1. Introduction.** Let  $F$  be a distribution function on  $[0, 1]^d$  and let  $X_i$ ,  $i = 1, 2, \dots$ , be independent random vectors taking values in  $[0, 1]^d$  with distribution function  $F$ . If one forms the empirical distribution function

$$F_n(\mathbf{t}) = \frac{1}{n} \sum_{i=1}^n 1_{[0, \mathbf{t}]}(X_i),$$

it is well known that  $\sqrt{n}(F_n(\cdot) - F(\cdot))$  converges in law (with respect to the supremum norm  $\|\cdot\|$ ) to the mean 0 Gaussian process  $W_F$ , where

$$\text{Cov}(W_F(\mathbf{s}), W_F(\mathbf{t})) = F([\mathbf{0}, \mathbf{s}] \cap [\mathbf{0}, \mathbf{t}]) - F([\mathbf{0}, \mathbf{s}])F([\mathbf{0}, \mathbf{t}]).$$

We refer to  $W_F$  as the tied-down Brownian process determined by  $F$ . Let  $G$  be a function on  $[0, 1]^d$ . The main purpose of this paper is to obtain upper and lower bounds for  $P(\|W_F - G\| \leq \epsilon)$ . We also obtain analogous bounds for  $P(\|B - G\| \leq \epsilon)$ , where  $B$  is the standard Brownian sheet.

The principal motivation is as follows. Given samples  $X_i$  as above, one wants to test whether they could have law  $F$ . One forms  $F_n$  as above, one defines the Kolmogorov–Smirnov statistic  $D_n = \sqrt{n}(F_n - F)$ , and one would like to reject the hypothesis that the  $X_i$ 's have law  $F$  if  $\|D_n\|$  is too large. The distribution of  $D_n$  cannot be calculated exactly, so instead one looks at  $\|W_F\|$ . Even here, in the case that the dimension  $d > 1$ , the distribution of the tail of  $W_F$  is not known, although successively better estimates have been obtained by Goodman (1976), Cabaña and Wschebor (1982) and Adler and Brown (1986). Now suppose one wants to determine the power of this test. [Blum, Kiefer and Rosenblatt (1961) have argued that tests of this type should have very good power.] This means one wants to estimate  $P(\|D_n\| \leq \alpha)$  when the  $X_i$ 's are in fact from another distribution  $H$ . Write

$$\begin{aligned} P(\|\sqrt{n}(F_n - F)\| \leq \alpha) &= P(\|\sqrt{n}(F_n - H) + \sqrt{n}(H - F)\| \leq \alpha) \\ &\approx P(\|W_H - K\| \leq \alpha), \end{aligned}$$

---

Received April 1986; revised December 1986.

<sup>1</sup>Partially supported by NSF Grant MCS-83-00581.

AMS 1980 subject classifications. Primary 60G15; secondary 60F10, 60G60, 62G10.

Key words and phrases. Brownian sheet, Kolmogorov–Smirnov, large deviations, Haar functions, empirical processes.

where  $K = \sqrt{n}(F - H)$ . Consequently, the estimates of this paper (Theorems 3.4 and 4.5) will give information on the power of the Kolmogorov–Smirnov test. Our bounds are clearly not sharp, and hopefully future research will refine them.

Our bounds may also be viewed as being of large deviation type. Using standard techniques, one can recover known results on the probability of large deviations for the Brownian sheet and the tied-down Brownian process [cf. Varadhan (1984), Theorem 5.1, for the one-parameter case]. Moreover, in a very rough sense, these estimates give some information as to what the second-order term should look like in Sanov-type inequalities [see Sanov (1957), Borovkov (1967) and Groeneboom, Oosterhoff and Ruymgaart (1979)]. Of course, there one really wants estimates for  $\|D_n - G\|$  and not  $\|W_F - G\|$ .

A third application of our estimates is related to the extension of Chung’s law of the iterated logarithm to the Brownian sheet. Our estimates can be easily used to show that if there exists a function  $\varphi(t)$  such that

$$\liminf_{t \rightarrow \infty} \sup_{s \in [0, (t, \dots, t)]} |B(\mathbf{s})| / \varphi(t)$$

equals a constant, a.s., then  $\varphi(t)$  must lie between  $t^{d/2}(\log \log \log t)^{(d-1)/2} / (\log \log t)^{1/2}$  and  $t^{d/2}(\log \log \log t)^{3(d-1)/2} / (\log \log t)^{1/2}$ . This improves a result of Révész (1981) who showed in the case  $d = 2$  that  $\varphi(t)$  must lie between  $t(\log \log \log t)^{1/2} / (\log \log t)^{1/2}$  and  $t(\log \log \log t)^{5/2} / (\log \log t)^{1/2}$ . The presence of the  $\log \log \log t$  is rather surprising, and it would be an interesting but quite difficult problem to determine if an exact normalizing function  $\varphi(t)$  exists, if it can be expressed in terms of an exact power of  $\log \log \log t$ , and if so, what power.

We obtain the upper bound by reducing the problem for the tied-down Brownian process determined by  $F$  to the corresponding problem for the untied process determined by  $F$ , then using stochastic integrals to reduce to the case of Brownian sheet, and finally using the Haar function decomposition of the Brownian sheet. This use of stochastic integrals is, as far as we know, new. The lower bound proceeds by the same sequence of reductions and also uses the Haar function representation, but, in addition, we use the Cameron–Martin–Girsanov transformation. Section 2 contains the necessary preliminaries concerning Haar functions and stochastic integrals, Section 3 contains the derivation of the upper bound, and Section 4 contains the derivation of the lower bound.

To simplify the notation, throughout we consider only the case  $d = 2$ , pointing out the changes necessary for  $d > 2$  when they are not straightforward. We say  $F: [0, 1]^d \rightarrow \mathbb{R}$  has a density  $f$  if  $F(\mathbf{t}) = \int_{[0, \mathbf{t}]} f(\mathbf{s}) d\mathbf{s}$  for all  $\mathbf{t}$ . The letter  $c$ , with or without subscripts, denotes constants whose value may change from line to line.

**2. Preliminaries.** Let  $F$  be a continuous distribution function on  $[0, 1]^2$ . We may, of course, view  $F$  as either a function or a measure. Define the Brownian sheet determined by  $F$ ,  $B_F$ , to be the mean 0 Gaussian process with continuous paths and covariance given by  $\text{Cov}(B_F(\mathbf{s}), B_F(\mathbf{t})) = F([\mathbf{0}, \mathbf{s}] \cap [\mathbf{0}, \mathbf{t}])$ . Define the tied-down Brownian process determined by  $F$ ,  $W_F$ , to be the mean 0 Gaussian process with continuous paths and covariance given by

$\text{Cov}(W_F(\mathbf{s}), W_F(\mathbf{t})) = F([\mathbf{0}, \mathbf{s}] \cap [\mathbf{0}, \mathbf{t}]) - F([\mathbf{0}, \mathbf{s}])F([\mathbf{0}, \mathbf{t}])$ . When  $F$  is the uniform distribution, i.e., when  $F([\mathbf{0}, \mathbf{s}]) = \|\mathbf{0}, \mathbf{s}\|$ , where  $|\cdot|$  denotes Lebesgue measure, we write simply  $B$  for the standard Brownian sheet.

Next, we recall the definition of the Haar functions. For  $m = 1, 2, \dots$ , let  $\Gamma(m) = \{1, 2, \dots, 2^{m-1}\}$ . Let  $\Gamma(0) = \{0\}$ . For  $m \geq 1$ ,  $j \in \Gamma(m)$ , define  $\phi_{jm}: [0, 1] \rightarrow \mathbb{R}$  by

$$\phi_{jm}(t) = \begin{cases} 2^{(m-1)/2}, & t \in \left[ \frac{2j-2}{2^m}, \frac{2j-1}{2^m} \right), \\ -2^{(m-1)/2}, & t \in \left[ \frac{2j-1}{2^m}, \frac{2j}{2^m} \right), \\ 0, & \text{otherwise.} \end{cases}$$

For  $j = 0, m = 0$ , let  $\phi_{jm}(t) = 1$  for  $t \in [0, 1]$ . It is well known that the Haar functions  $\phi_{jm}$  form a complete orthonormal system for  $L^2([0, 1], dt)$ .

Now if  $\mathbf{m} = (m_1, m_2)$ , let  $\Gamma(\mathbf{m}) = \{(j_1, j_2): j_1 \in \Gamma(m_1), j_2 \in \Gamma(m_2)\}$ . If  $\mathbf{j} = (j_1, j_2) \in \Gamma(\mathbf{m})$  and  $\mathbf{t} = (t_1, t_2) \in [0, 1]^2$ , define

$$\Phi_{\mathbf{j}\mathbf{m}}(\mathbf{t}) = \phi_{j_1, m_1}(t_1)\phi_{j_2, m_2}(t_2).$$

The  $\Phi_{\mathbf{j}\mathbf{m}}$  form a complete orthonormal system for  $L^2([0, 1]^2, dt)$ .

Let  $\psi(\mathbf{m})$  be defined so that the cardinality of  $\Gamma(\mathbf{m})$  is  $2^{\psi(\mathbf{m})}$ . Thus

$$\psi(\mathbf{m}) = (m_1 - 1)1_{(m_1 > 0)} + (m_2 - 1)1_{(m_2 > 0)}.$$

Also, observe that the number of  $\mathbf{m}$ 's for which  $\psi(\mathbf{m}) = k$  is  $k + 3$  for  $k > 0$ , 4 if  $k = 0$ .

Let  $\langle f, g \rangle$  denote  $\int_{[0, 1]^2} f(\mathbf{t})g(\mathbf{t}) d\mathbf{t}$ , and let  $\|f\|$  denote  $\sup_{\mathbf{t} \in [0, 1]^2} |f(\mathbf{t})|$ . Let

$$\alpha_{\mathbf{j}\mathbf{m}}(\mathbf{t}) = \langle 1_{[0, \mathbf{t}]} , \Phi_{\mathbf{j}\mathbf{m}} \rangle.$$

A simple calculation shows that  $\|\alpha_{\mathbf{j}\mathbf{m}}\| \leq 2(2^{-\psi(\mathbf{m})/2})$ , while  $\|\Phi_{\mathbf{j}\mathbf{m}}\| = 2^{\psi(\mathbf{m})/2}$  is immediate from the definitions.

The support of  $t \rightarrow \int_0^{t_1} \phi_{j_1, m_1}(s) ds$  is the same as the support of  $\phi_{j_1, m_1}$ . Since by Fubini,

$$\alpha_{\mathbf{j}\mathbf{m}}(\mathbf{t}) = \left( \int_0^{t_1} \phi_{j_1, m_1}(s_1) ds_1 \right) \left( \int_0^{t_2} \phi_{j_2, m_2}(s_2) ds_2 \right),$$

it follows that the support of  $\alpha_{\mathbf{j}\mathbf{m}}$  is the same as the support of  $\Phi_{\mathbf{j}\mathbf{m}}$ . For each  $\mathbf{m}$ , the functions  $\Phi_{\mathbf{j}\mathbf{m}}, \mathbf{j} \in \Gamma(\mathbf{m})$ , have disjoint support, and hence for each  $\mathbf{m}$ , each point  $\mathbf{t}$  is in the support of at most one  $\alpha_{\mathbf{j}\mathbf{m}}$ . So for fixed  $\mathbf{m}$  and constants  $d_{\mathbf{j}\mathbf{m}}$ , we have

$$(2.1) \quad \left\| \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} d_{\mathbf{j}\mathbf{m}} \alpha_{\mathbf{j}\mathbf{m}} \right\| \leq \sup_{\mathbf{j} \in \Gamma(\mathbf{m})} |d_{\mathbf{j}\mathbf{m}}| \|\alpha_{\mathbf{j}\mathbf{m}}\|.$$

We also will need Parseval's identities,

$$(2.2) \quad \sum_{\mathbf{j}, \mathbf{m}} \langle f, \Phi_{\mathbf{j}\mathbf{m}} \rangle^2 = \langle f, f \rangle$$

and

$$(2.3) \quad \sum_{\mathbf{j}, \mathbf{m}} \langle f, \Phi_{\mathbf{j}\mathbf{m}} \rangle \langle g, \Phi_{\mathbf{j}\mathbf{m}} \rangle = \langle f, g \rangle,$$

where the sum is over all  $\mathbf{m}$  and all  $\mathbf{j} \in \Gamma(\mathbf{m})$ .

We now recall the construction of Brownian sheet by means of Haar functions [see Park (1970)]. Let  $Z_{\mathbf{j}\mathbf{m}}$  be independent standard normal variables. For each  $\mathbf{t}$ ,  $N \geq 1$ , define

$$(2.4) \quad B_N(\mathbf{t}) = \sum_{\{\mathbf{m}: \psi(\mathbf{m}) \leq N\}} \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} Z_{\mathbf{j}\mathbf{m}} \alpha_{\mathbf{j}\mathbf{m}}(\mathbf{t}).$$

Clearly,  $B_N(\mathbf{t})$  is a mean 0 Gaussian process with continuous paths. It is known that  $\|B_N - B\| \rightarrow 0$  in  $L^2(\Omega, P)$ , where  $B$  is the process defined by

$$(2.5) \quad B(\mathbf{t}) = \sum_{\mathbf{m}} \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} Z_{\mathbf{j}\mathbf{m}} \alpha_{\mathbf{j}\mathbf{m}}(\mathbf{t}).$$

The fact that  $B$  is a mean 0 Gaussian process with continuous paths follows from the corresponding fact about  $B_N$ . Using (2.3), we have

$$\begin{aligned} \text{Cov}(B(\mathbf{s}), B(\mathbf{t})) &= \sum_{\mathbf{m}} \sum_{\mathbf{j}} \alpha_{\mathbf{j}\mathbf{m}}(\mathbf{t}) \alpha_{\mathbf{j}\mathbf{m}}(\mathbf{s}) = \sum_{\mathbf{m}, \mathbf{j}} \langle 1_{[0, \mathbf{t}]}, \Phi_{\mathbf{j}\mathbf{m}} \rangle \langle 1_{[0, \mathbf{s}]}, \Phi_{\mathbf{j}\mathbf{m}} \rangle \\ &= \langle 1_{[0, \mathbf{t}]}, 1_{[0, \mathbf{s}]} \rangle = |[0, \mathbf{t}] \cap [0, \mathbf{s}]|. \end{aligned}$$

We next turn to stochastic integrals. If  $h(\mathbf{t}) = \sum_{i=1}^I d_i 1_{[0, \mathbf{r}_i]}(\mathbf{t})$  is simple, the stochastic integral  $\int h dB$  is defined to be  $\sum_{i=1}^I d_i B(\mathbf{r}_i)$ . Then  $\int h dB$  is mean 0, Gaussian,  $\mathbf{t} \rightarrow \int 1_{[0, \mathbf{t}]} h dB$  has continuous paths, and using the independent increments property of the Brownian sheet, an easy calculation gives

$$(2.6) \quad E \left( \int h dB \right)^2 = \int h^2(\mathbf{t}) d\mathbf{t}.$$

Applying (2.6) to  $h, k$  and  $h + k$ , we get by polarization that

$$(2.7) \quad E \left( \int h dB \right) \left( \int k dB \right) = \int h(\mathbf{t}) k(\mathbf{t}) d\mathbf{t}.$$

An argument using Doob's inequality applied twice [cf. Cairoli (1970)] shows that if  $Y(\mathbf{t}) = \int 1_{[0, \mathbf{t}]} h dB$ , then

$$(2.8) \quad E \|Y\|^2 \leq 16 \int h(\mathbf{t})^2 d\mathbf{t}.$$

For  $h \in L^2([0, 1]^2, d\mathbf{t})$ , choose  $h_n$  simple converging in  $L^2$  to  $h$ , and define  $\int h dB$  as the  $L^2(\Omega, dP)$  limit of  $\int h_n dB$ . The equality (2.6) is used to show that the limit exists and is independent of the choice of  $h_n$ . Again,  $\int h dB$  is mean 0, Gaussian, and (2.6) and (2.7) hold. Just as in the one-parameter case, (2.8) is used to show that  $\mathbf{t} \rightarrow \int 1_{[0, \mathbf{t}]} h dB$  has continuous paths a.s.

If  $F$  is a distribution function with a density  $f$ , then the process  $Y(\mathbf{t}) = \int 1_{[0, \mathbf{t}]} \sqrt{f} dB$  is a mean 0 continuous Gaussian process. Applying (2.7) with  $h = 1_{[0, \mathbf{s}]} \sqrt{f}$ ,  $k = 1_{[0, \mathbf{t}]} \sqrt{f}$ , we get

$$\text{Cov}(Y(\mathbf{s}), Y(\mathbf{t})) = \int 1_{[0, \mathbf{s}]} 1_{[0, \mathbf{t}]} f = F([0, \mathbf{s}] \cap [0, \mathbf{t}]).$$

Hence  $Y(\mathbf{t})$  has the same law as  $B_F(\mathbf{t})$ .

We need the following simple properties of stochastic integrals.

**LEMMA 2.1.** *If  $h \in L^2([0, 1]^2, d\mathbf{t})$ , then  $E(\int h dB_N - \int h dB)^2 \rightarrow 0$  as  $N \rightarrow \infty$ .*

**PROOF.** Note  $B_N(\mathbf{t}) = \int_{[0, \mathbf{t}]} b_N(\mathbf{s}) d\mathbf{s}$ , where

$$(2.9) \quad b_N(\mathbf{s}) = \sum_{\{\mathbf{m}: \psi(\mathbf{m}) \leq N\}} \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} Z_{\mathbf{j}\mathbf{m}} \Phi_{\mathbf{j}\mathbf{m}}(\mathbf{s}).$$

Hence the first integral in the statement of the lemma makes sense. For any  $H \in L^2$ ,  $\int H dB_N = \int H b_N d\mathbf{s}$  is mean 0, and recalling (2.2),

$$(2.10) \quad \begin{aligned} E\left(\int H b_N d\mathbf{s}\right)^2 &= \sum_{\{\mathbf{m}: \psi(\mathbf{m}) \leq N\}} \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} \left(\int H \Phi_{\mathbf{j}\mathbf{m}} d\mathbf{s}\right)^2 \\ &\leq \sum_{\mathbf{m}, \mathbf{j}} \langle H, \Phi_{\mathbf{j}\mathbf{m}} \rangle^2 = \langle H, H \rangle. \end{aligned}$$

Next, if  $K = \sum d_i 1_{[0, \mathbf{r}_i]}$  is simple,

$$\int K dB_N = \sum d_i B_N(\mathbf{r}_i) \rightarrow \sum d_i B(\mathbf{r}_i) = \int K dB,$$

as  $N \rightarrow \infty$ , the convergence being in  $L^2(\Omega, dP)$ .

Given  $\epsilon$ , choose  $K$  simple so that  $\langle K - h, K - h \rangle \leq \epsilon$ . Let  $H = K - h$ . Then by the triangle inequality, (2.6), and (2.10),

$$\begin{aligned} \limsup_{N \rightarrow \infty} \left( E\left(\int h dB_N - \int h dB\right)^2 \right)^{1/2} \\ \leq 2\epsilon + \limsup_{N \rightarrow \infty} \left( E\left(\int K dB_N - \int K dB\right)^2 \right)^{1/2} = 2\epsilon. \quad \square \end{aligned}$$

**LEMMA 2.2.** *Suppose  $M$  has density  $m$  in  $L^2([0, 1]^2, d\mathbf{t})$  and  $H$  has density  $h$  in  $L^1([0, 1]^2, d\mathbf{t})$ . Suppose  $K \in L^2([0, 1]^2, d\mathbf{t})$ . Then*

$$\left\| \int_{[0, \mathbf{t}]} HK dB - \int_{[0, \mathbf{t}]} HK dM \right\| \leq 4 \left( \int |h(\mathbf{r})| d\mathbf{r} \right) \left\| \int_{[0, \cdot]} K dB - \int_{[0, \cdot]} K dM \right\|.$$

PROOF. It clearly suffices to take  $h$  bounded, the general case following by taking a limit. Let

$$Y_N(\mathbf{t}) = \int_{[0, \mathbf{t}]} K d(B_N - M), \quad Y(\mathbf{t}) = \int_{[0, \mathbf{t}]} K dB - \int_{[0, \mathbf{t}]} K dM.$$

By Lemma 2.1,  $E(Y_N(\mathbf{t}) - Y(\mathbf{t}))^2 \rightarrow 0$  for each  $\mathbf{t}$ , and moreover, by (2.10),  $E(Y_N(\mathbf{t}) - Y(\mathbf{t}))^2 \leq 4\langle K, K \rangle$ . So by dominated convergence,

$$(2.11) \quad E \left| \int [Y_N(\mathbf{t}) - Y(\mathbf{t})] h(\mathbf{t}) d\mathbf{t} \right| \leq \|h\| \int (E(Y_N(\mathbf{t}) - Y(\mathbf{t}))^2)^{1/2} d\mathbf{t} \rightarrow 0.$$

Recalling the definition of  $b_N(\mathbf{s})$  in (2.9), we have

$$(2.12) \quad \begin{aligned} & \left| \int_{[0, \mathbf{t}]} H(\mathbf{s}) K(\mathbf{s}) [b_N(\mathbf{s}) - m(\mathbf{s})] d\mathbf{s} \right| \\ &= \left| \int_{[0, \mathbf{t}]} K(\mathbf{s}) [b_N(\mathbf{s}) - m(\mathbf{s})] \int_{[0, \mathbf{s}]} h(\mathbf{r}) d\mathbf{r} d\mathbf{s} \right| \\ &= \left| \int_{[0, \mathbf{t}]} h(\mathbf{r}) \int_{[\mathbf{r}, \mathbf{t}]} K(\mathbf{s}) [b_N(\mathbf{s}) - m(\mathbf{s})] d\mathbf{s} d\mathbf{r} \right| \\ &= \left| \int [Y_N(\mathbf{t}) + Y_N(\mathbf{r}) - Y_N(r_1, t_2) - Y_N(t_1, r_2)] h(\mathbf{r}) d\mathbf{r} \right|. \end{aligned}$$

Let  $N \rightarrow \infty$ . Then  $\int Y_N(\mathbf{r}) h(\mathbf{r}) d\mathbf{r} \rightarrow \int Y(\mathbf{r}) h(\mathbf{r}) d\mathbf{r}$  in  $L^1(\Omega, dP)$  by (2.11). The other terms may be treated similarly, and so the right-hand side of (2.12) converges in  $L^1(\Omega, dP)$  to

$$\left| \int [Y(\mathbf{t}) + Y(\mathbf{r}) - Y(r_1, t_2) - Y(t_1, r_2)] h(\mathbf{r}) d\mathbf{r} \right| \leq 4\|Y\| \int |h(\mathbf{r})| d\mathbf{r}.$$

By Lemma 2.1, the left-hand side of (2.12) converges to  $|\int_{[0, \mathbf{t}]} HK dB - \int_{[0, \mathbf{t}]} HK dM|$ , which completes the proof.  $\square$

Finally, we need to recall the Cameron–Martin–Girsanov theorem. There are several proofs in the literature, but using the Haar decomposition of the Brownian sheet, we can give the following simple proof.

THEOREM 2.3. Suppose  $h \in L^2([0, 1]^2, dt)$ . Define  $Q$  by

$$dQ/dP = \exp\left(\int h dB - \frac{1}{2} \int h^2 dt\right).$$

Then under  $Q$ ,  $B$  is a Gaussian process with  $\text{Cov}_Q(B(\mathbf{s}), B(\mathbf{t})) = |[0, \mathbf{s}] \cap [0, \mathbf{t}]|$  and  $E_Q B(\mathbf{t}) = \int_{[0, \mathbf{t}]} h$ .

PROOF. Write  $h$  as  $\sum_{j,m} \langle h, \Phi_{jm} \rangle \Phi_{jm}$ . Then  $\int h dB = \sum_{j,m} \langle h, \Phi_{jm} \rangle Z_{jm}$ . Recalling the independence of the  $Z_{jm}$ 's, (2.2) and (2.3),

$$\begin{aligned} E_Q \exp(iuB(t)) &= E_P \exp\left(\sum_{j,m} (iu\langle 1_{[0,t]}, \Phi_{jm} \rangle + \langle h, \Phi_{jm} \rangle) Z_{jm}\right) \exp\left(-\frac{1}{2} \int h^2\right) \\ &= \prod_{j,m} \exp\left(\frac{1}{2} (iu\langle 1_{[0,t]}, \Phi_{jm} \rangle + \langle h, \Phi_{jm} \rangle)^2\right) \exp\left(-\frac{1}{2} \int h^2\right) \\ &= \exp\left(-\frac{1}{2} u^2 \sum_{j,m} \langle 1_{[0,t]}, \Phi_{jm} \rangle^2 + iu \sum_{j,m} \langle 1_{[0,t]}, \Phi_{jm} \rangle \langle h, \Phi_{jm} \rangle \right. \\ &\qquad \qquad \qquad \left. + \frac{1}{2} \sum_{j,m} \langle h, \Phi_{jm} \rangle^2\right) \exp\left(-\frac{1}{2} \int h^2\right) \\ &= \exp\left(-\frac{1}{2} u^2 \langle 1_{[0,t]}, 1_{[0,t]} \rangle + iu \langle 1_{[0,t]}, h \rangle\right). \end{aligned}$$

So under  $Q$ ,  $B(t)$  is Gaussian with mean  $\int h 1_{[0,t]}$  and variance  $[[0, t]]$ . A similar calculation for  $E_Q \exp(i \sum_k u_k B(t_k))$  shows that the finite-dimensional distributions of  $B$  under  $Q$  are Gaussian and that the covariances under  $Q$  are the same as those under  $P$ .  $\square$

**3. Upper bound.** We proceed by a series of reductions.

PROPOSITION 3.1. *Suppose  $G$  is continuous and  $G(1) = 0$ . Then*

$$P(\|W_F - G\| \leq \epsilon) \leq c(\epsilon) P(\|B_F - G\| \leq 2\epsilon),$$

where  $c(\epsilon) = O(1/\epsilon)$  as  $\epsilon \rightarrow 0$ .

PROOF. A standard calculation involving conditional distributions shows that the distribution of  $B_F(\cdot)$  given  $B_F(1) = 0$  is that of a mean 0 Gaussian process with the same covariance structure as  $W_F(\cdot)$ . The distribution of  $B_F(\cdot)$  given  $B_F(1) = x$  has the same covariance structure, and the mean is readily calculated:

$$\begin{aligned} E(B_F(t) | B_F(1) = x) &= E(B_F(t) - F(t)B_F(1) | B_F(1) = x) + F(t)x \\ &= E(B_F(t) - F(t)B_F(1)) + F(t)x = F(t)x, \end{aligned}$$

where in the next to the last equality we used the fact that  $B_F(t) - F(t)B_F(1)$  and  $B_F(1)$  are Gaussian random variables whose covariance is 0, hence they are independent.

We then have

$$\begin{aligned} P(\|B_F - G\| \leq 2\epsilon) &\geq P(\|B_F - G\| \leq 2\epsilon, |B_F(1)| \leq \epsilon) \\ &= \int_{-\epsilon}^{\epsilon} P(\|B_F - G\| \leq 2\epsilon | B_F(1) = x) P(B_F(1) \in dx) \\ &\geq \int_{-\epsilon}^{\epsilon} P(\|B_F - F(\cdot)x - G\| \leq \epsilon | B_F(1) = x) P(B_F(1) \in dx) \\ &\geq (c(\epsilon))^{-1} P(\|W_F - G\| \leq \epsilon). \quad \square \end{aligned}$$

PROPOSITION 3.2. Suppose  $G(1) = 0$ ,  $G$  has a density  $g$  in  $L^2([0, 1]^2, dt)$ , and  $F$  has a density  $f$  with

$$\frac{d(f^{1/2})}{dt}, \frac{d(f^{-1/2})}{dt} \in L^1([0, 1]^2, dt).$$

Then

$$P(\|B_F - G\| \leq \epsilon) \leq P(\|B - L\| \leq c(F)\epsilon),$$

where  $L$  is the function with density  $g/\sqrt{f}$ .

PROOF. By Section 2,  $\int_{[0,t]} f^{1/2} dB$  has the same law as  $B_F$ . By Lemma 2.2 (with  $H = f^{-1/2}$ ,  $K = f^{1/2}$ ,  $M = L$ ),

$$\|B(t) - L(t)\| \leq (c(F))^{-1} \left\| \int_{[0, \cdot]} f^{1/2} dB - \int_{[0, \cdot]} f^{1/2} dL \right\| = (c(F))^{-1} \|B_F - G\|.$$

The proposition is now immediate.  $\square$

PROPOSITION 3.3. If  $H$  has a density  $h \in L^2([0, 1]^2, dt)$ , then

$$P(\|B - H\| \leq \epsilon) \leq c_1 \exp\left(-\frac{1}{2} \int_{[0,1]^2} h^2 dt\right) \exp(-c_2 \epsilon^{-2} \ln(1/\epsilon)).$$

PROOF. By the definition of stochastic integrals,

$$\begin{aligned} & \left| \int 1_{[r,s]} dB - \int 1_{[r,s]} dH \right| \\ &= |(B - H)(s) + (B - H)(r) - (B - H)(r_1, s_2) - (B - H)(s_1, r_2)| \\ &\leq 4\|B - H\|. \end{aligned}$$

Since any Haar function  $\Phi_{jm}$  can be written as  $\sum_i d_{jm}^{(i)} 1_{[r_i, s_i]}$ , where the sum is from  $i = 1$  to  $i = 1, 2$  or  $4$  (depending on  $\mathbf{j}, \mathbf{m}$ ) and  $d_{jm}^{(i)} = \pm \|\Phi_{jm}\|$ , then  $|\int \Phi_{jm} d(B - H)| \leq 16\|\Phi_{jm}\| \|B - H\|$ . On the other hand,  $\int \Phi_{jm} d(B - H) = Z_{jm} - \langle \Phi_{jm}, h \rangle$ . So if  $\|B - H\| \leq \epsilon$ , then for each  $\mathbf{j}$  and  $\mathbf{m}$ ,  $|Z_{jm} - \langle \Phi_{jm}, h \rangle| \leq 16(2^{\psi(\mathbf{m})/2})\epsilon$ . Since the  $Z_{jm}$  are independent, then for each  $M$ ,

$$(3.1) \quad P(\|B - H\| \leq \epsilon) \leq \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq M\}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} p_{jm},$$

where

$$p_{jm} = P(|Z_{jm} - \langle \Phi_{jm}, h \rangle| \leq 16(2^{\psi(\mathbf{m})/2})\epsilon).$$

By the trivial inequality

$$P(Z \in A) \leq (2\pi)^{-1/2} \left( \sup_{y \in A} e^{-y^2/2} \right) |A|,$$

for  $Z$  a standard normal random variable, we have that for each  $\mathbf{j}$  and  $\mathbf{m}$ ,

$$(3.2) \quad p_{jm} \leq (2\pi)^{-1/2} 16(2^{\psi(\mathbf{m})/2})\epsilon.$$

By this same inequality, we have

$$(3.3) \quad \begin{aligned} P_{\mathbf{j}\mathbf{m}} &\leq (2\pi)^{-1/2} 16(2^{\psi(\mathbf{m})/2})\varepsilon \exp\left(-\frac{1}{2}\left(|\langle \Phi_{\mathbf{j}\mathbf{m}}, h \rangle| - 16(2^{\psi(\mathbf{m})/2})\varepsilon\right)^2\right) \\ &\leq 16(2\pi)^{-1/2} 2^{\psi(\mathbf{m})/2}\varepsilon \exp\left(-\frac{1}{2}\langle \Phi_{\mathbf{j}\mathbf{m}}, h \rangle^2\right) \exp(c(\mathbf{j}, \mathbf{m}, h)(\varepsilon + o(\varepsilon))) \end{aligned}$$

provided  $|\langle \Phi_{\mathbf{j}\mathbf{m}}, h \rangle| > 16(2^{\psi(\mathbf{m})/2})\varepsilon$ .

Now let  $\delta \in (0, 1)$  be fixed, and choose  $K$  large so that

$$\sum_{\{\mathbf{m}: \psi(\mathbf{m}) \leq K\}} \sum_{\mathbf{j}} \langle h, \Phi_{\mathbf{j}\mathbf{m}} \rangle^2 \geq \langle h, h \rangle - \delta.$$

Then take  $\varepsilon$  small enough so that for every  $\mathbf{m}$  with  $\psi(\mathbf{m}) \leq K$  and every  $\mathbf{j} \in \Gamma(\mathbf{m})$ , either  $|\langle \Phi_{\mathbf{j}\mathbf{m}}, h \rangle| = 0$  or  $> 16(2^{\psi(\mathbf{m})/2})\varepsilon$ . Pick  $M$  so that  $16(2\pi)^{-1/2}(2^{M/2}) \in [\varepsilon^{-1}/16, \varepsilon^{-1}]$ . If necessary, choose  $\varepsilon$  even smaller so that  $M > K$ .

Observe by our choice of  $\varepsilon$  and (3.2) that (3.3) holds for every  $\mathbf{m}$  such that  $\psi(\mathbf{m}) \leq K$  and  $\mathbf{j} \in \Gamma(\mathbf{m})$ . Since  $K$  does not depend on  $\varepsilon$ , we have

$$(3.4) \quad \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq K\}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} \exp(c(\mathbf{j}, \mathbf{m}, h)(\varepsilon + o(\varepsilon))) = O(1),$$

as  $\varepsilon \rightarrow 0$ .

Using (3.3) when  $\psi(\mathbf{m}) \leq K$  and (3.2) when  $K < \psi(\mathbf{m}) \leq M$ , substituting into (3.1), and recalling (3.4), we have

$$(3.5) \quad \begin{aligned} P(\|B - H\| \leq \varepsilon) &\leq c_1 \left( \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq K\}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} \exp\left(-\frac{1}{2}\langle \Phi_{\mathbf{j}\mathbf{m}}, h \rangle^2\right) \right) \\ &\quad \times \left( \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq M\}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} 16(2\pi)^{-1/2} 2^{\psi(\mathbf{m})/2}\varepsilon \right). \end{aligned}$$

By the choice of  $K$ , the product of the exponentials is

$$\leq \exp\left(-\frac{1}{2}(\langle h, h \rangle - \delta)\right) \leq c_3 \exp\left(-\frac{1}{2}\langle h, h \rangle\right),$$

and so to complete the proof, it suffices to bound

$$\begin{aligned} q &= \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq M\}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} 16(2\pi)^{-1/2} 2^{\psi(\mathbf{m})/2}\varepsilon \\ &= \prod_{\{\mathbf{m}: \psi(\mathbf{m}) \leq M\}} (c2^{\psi(\mathbf{m})/2}\varepsilon) 2^{\psi(\mathbf{m})}, \end{aligned}$$

where by the choice of  $M$ ,  $c2^{\psi(\mathbf{m})/2}\varepsilon \leq 1$ . Taking logarithms, we get

$$(3.6) \quad \begin{aligned} \ln q &\leq \sum_{k=0}^M \sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} 2^k \ln(c2^{k/2}\varepsilon) \\ &\leq \sum_{k=0}^M k2^k \ln(c2^{k/2}\varepsilon) \\ &\leq c_1 \int_1^{2^M} \ln x \ln(c_2 x \varepsilon^2) dx \\ &= c_1 \varepsilon^{-2} \int_{\varepsilon^2}^{2^M \varepsilon^2} \ln y \ln(c_2 y) dy + 2c_1 \varepsilon^{-2} \ln(1/\varepsilon) \int_{\varepsilon^2}^{2^M \varepsilon^2} \ln(c_2 y) dy \\ &\leq -c_3 \varepsilon^{-2} \ln(1/\varepsilon), \end{aligned}$$

if  $\epsilon \leq 1/256$ . To get the last inequality, we used the facts that

$$\int_{\epsilon^2}^{2^M \epsilon^2} \geq \int_{1/(256)^2}^{2\pi/(256)^2}$$

and that the integrands do not depend on  $\epsilon$ .  $\square$

**REMARK.** For  $d > 2$ ,  $\sum_{k=0}^M \sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} 1 \geq \sum_{k=0}^M k^{d-1}$ , provided  $\psi(\mathbf{m})$  is defined in the obvious way. With this change in (3.6), we obtain  $-c_3 \epsilon^{-2} (\ln(1/\epsilon))^{d-1}$ .

**THEOREM 3.4.** Suppose  $G(\mathbf{1}) = 0$ ,  $G$  has a density  $g$ ,  $F$  has a density  $f$ ,

$$\frac{d(f^{1/2})}{dt}, \frac{d(f^{-1/2})}{dt} \in L^1([0, 1]^d, dt) \quad \text{and} \quad g/\sqrt{f} \in L^2([0, 1]^d, dt).$$

Then

$$P(\|W_F - G\| \leq \epsilon) \leq c_1 \exp\left(-\frac{1}{2} \int_{[0, 1]^d} (g^2/f) dt - c_2 \epsilon^{-2} (\ln(1/\epsilon))^{d-1}\right),$$

where  $c_1$  and  $c_2$  are constants depending only on  $F$ .

**PROOF.** Combine Propositions 3.1, 3.2 and 3.3.  $\square$

**4. Lower bound.** Again, we proceed by a series of reductions.

**PROPOSITION 4.1.** If  $G(\mathbf{1}) = 0$ , then

$$P(\|W_F - G\| \leq \epsilon) \geq P(\|B_F - G\| \leq \epsilon/2).$$

**PROOF.** Define  $Y(\mathbf{t}) = B_F(\mathbf{t}) - F(\mathbf{t})B_F(\mathbf{1})$ . Clearly,  $Y$  is mean 0 Gaussian with continuous paths, and a simple calculation shows that the covariances of  $Y$  are the same as those of  $W_F$ , hence the law of  $Y$  is the same as the law of  $W_F$ . But since  $\|F\| \leq 1$ , we have that if  $\|B_F - G\| \leq \epsilon/2$ , then in particular  $|B_F(\mathbf{1})| \leq \epsilon/2$ , and so  $\|Y - G\| \leq \epsilon$ . The proof is now immediate.  $\square$

**PROPOSITION 4.2.** Suppose  $F$  has a density  $f$  with  $d(f^{1/2})/dt \in L^1([0, 1]^2, dt)$ . Suppose  $G$  has a density  $g$  with  $g/\sqrt{f} \in L^2([0, 1]^2, dt)$ . Let  $L$  be the function with density  $g/\sqrt{f}$ . Then

$$P(\|B_F - G\| \leq \epsilon) \geq P(\|B - L\| \leq c\epsilon).$$

**PROOF.** Apply Lemma 2.2 (with  $K = 1$ ,  $M = L$  and  $H = f^{1/2}$ ) to get  $|B_F(\mathbf{t}) - G(\mathbf{t})| \leq c\|B - L\|$ .  $\square$

**PROPOSITION 4.3.** If  $H$  has a density  $h \in L^2([0, 1]^2, dt)$ , then

$$P(\|B - H\| \leq \epsilon) \geq \exp\left(-\frac{1}{2} \int h^2 dt\right) P(\|B\| \leq \epsilon).$$

PROOF [cf. Csáki (1980)]. By the symmetry of the Gaussian measure, for all  $\mathbf{r}$  and  $\mathbf{s}_1, \dots, \mathbf{s}_k$ ,

$$E(B(\mathbf{r}); |B(\mathbf{s}_1)| < \varepsilon, \dots, |B(\mathbf{s}_k)| < \varepsilon) = 0.$$

By taking limits, we get  $E(B(\mathbf{r}); \|B\| < \varepsilon) = 0$ . Using this, we have

$$(4.1) \quad E\left(\int h dB; \|B\| < \varepsilon\right) = 0,$$

if  $h$  is simple, and by taking limits, we have (4.1) for all  $h \in L^2$ .

Define a probability measure  $Q$  by  $dQ/dP = \exp(-\int h dB - \frac{1}{2}\int h^2 dt)$ . By Theorem 2.3, the law of  $B$  under  $Q$  is that of a Gaussian process,  $B$  has continuous paths under  $Q$  (since  $P$  and  $Q$  are equivalent),  $\text{Cov}_Q(B(\mathbf{s}), B(\mathbf{t})) = |\mathbf{0}, \mathbf{s}| \cap [0, \mathbf{t}]$  and  $E_Q B(\mathbf{t}) = -H(\mathbf{t})$ . Thus the law of  $B$  under  $Q$  is the same as the law of  $B - H$  under  $P$ . Therefore

$$(4.2) \quad P(\|B - H\| \leq \varepsilon) = Q(\|B\| \leq \varepsilon) = \int_{(\|B\| \leq \varepsilon)} \exp\left(-\int h dB - \frac{1}{2}\int h^2 dt\right) dP.$$

Using Jensen's inequality with respect to  $P(\cdot | \|B\| \leq \varepsilon)$ , we get

$$(4.3) \quad \int_{(\|B\| \leq \varepsilon)} \exp\left(-\int h dB\right) dP \geq P(\|B\| \leq \varepsilon) \exp\left(\int_{(\|B\| \leq \varepsilon)} \left(\int -h dB\right) dP / P(\|B\| \leq \varepsilon)\right).$$

Combining (4.1), (4.2) and (4.3) completes the proof.  $\square$

PROPOSITION 4.4.  $P(\|B\| \leq \varepsilon) \geq c_1 \exp(-c_2 \varepsilon^{-2} (\ln(1/\varepsilon))^3)$ .

PROOF. Fix  $\varepsilon$  small, and choose  $N$  so that  $2^N \varepsilon^2 / N^2 \in [1, 16]$ . Let  $\beta_{\mathbf{m}} = (3/\pi^2(\psi(\mathbf{m}) + 4)(N - \psi(\mathbf{m}) + 1)^{-2})$ , so that  $\sum_{\mathbf{m}} \beta_{\mathbf{m}} = \sum_{k=0}^{\infty} \sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} \beta_{\mathbf{m}} \leq 1$ . Let  $r_{\mathbf{j}\mathbf{m}} = \ln P(|Z_{\mathbf{j}\mathbf{m}}| < 2^{\psi(\mathbf{m})/2} \beta_{\mathbf{m}} \varepsilon / 2)$ .

By (2.1) with  $d_{\mathbf{j}\mathbf{m}} = Z_{\mathbf{j}\mathbf{m}}(\omega)$ , we have that if  $\sup_{\mathbf{j} \in \Gamma(\mathbf{m})} \|\alpha_{\mathbf{j}\mathbf{m}}\| |Z_{\mathbf{j}\mathbf{m}}| < \beta_{\mathbf{m}} \varepsilon$  for each  $\mathbf{m}$ , then

$$\|B\| \leq \sum_{\mathbf{m}} \left\| \sum_{\mathbf{j} \in \Gamma(\mathbf{m})} Z_{\mathbf{j}\mathbf{m}} \alpha_{\mathbf{j}\mathbf{m}} \right\| < \sum_{\mathbf{m}} \beta_{\mathbf{m}} \varepsilon \leq \varepsilon.$$

So using the independence of the  $Z_{\mathbf{j}\mathbf{m}}$ 's, we have

$$(4.4) \quad \begin{aligned} P(\|B\| \leq \varepsilon) &\geq \prod_{\mathbf{m}} P\left(\sup_{\mathbf{j} \in \Gamma(\mathbf{m})} \|\alpha_{\mathbf{j}\mathbf{m}}\| |Z_{\mathbf{j}\mathbf{m}}| < \beta_{\mathbf{m}} \varepsilon\right) \\ &= \prod_{\mathbf{m}} P\left(\sup_{\mathbf{j} \in \Gamma(\mathbf{m})} |Z_{\mathbf{j}\mathbf{m}}| < 2^{\psi(\mathbf{m})/2} \beta_{\mathbf{m}} \varepsilon / 2\right) \\ &= \prod_{\mathbf{m}} \prod_{\mathbf{j} \in \Gamma(\mathbf{m})} \exp(r_{\mathbf{j}\mathbf{m}}). \end{aligned}$$

We now proceed to estimate. Recalling that the  $r_{jm}$  are  $\leq 0$  we have

$$\begin{aligned}
 (4.5) \quad q_1 &= \sum_{\{\mathbf{m}: \psi(\mathbf{m}) \leq N\}} \sum_{j \in \Gamma(\mathbf{m})} r_{jm} \\
 &\geq \sum_{k=0}^N \sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} 2^k \ln P(|Z| < 2^{k/2} \beta_{\mathbf{m}} \epsilon / 2),
 \end{aligned}$$

where  $Z$  is a standard normal variable. By our choice of  $N$ ,  $2^{k/2} \beta_{\mathbf{m}} \epsilon / 2$  is bounded above and below by constants independent of  $N$  and  $\epsilon$ , hence

$$P(|Z| < 2^{k/2} \beta_{\mathbf{m}} \epsilon / 2) \geq c 2^{k/2} \beta_{\mathbf{m}} \epsilon / 2,$$

and so

$$\begin{aligned}
 q_1 &\geq \sum_{k=0}^N (k+4) 2^k \ln \left( \frac{c 2^{k/2} \epsilon}{(k+4)(N-k+1)^2} \right) \\
 &\geq c_1 \sum_{k=1}^N k 2^k \ln \left( \frac{c_2 2^k \epsilon^2}{k^2 (N+1-k)^4} \right) + o(\epsilon^{-2}) \\
 &\geq c_1 \int_2^{2^N} \ln x \ln \left( \frac{c_2 x \epsilon^2}{(\ln x)^4 (N+2-\ln x)^4} \right) dx + o(\epsilon^{-2}) \\
 &\geq c_1 \int_2^{2^N} \ln x \ln \left( \frac{c_2 x \epsilon^2}{(\ln x)^2} \right) dx \\
 &\quad - c_1 \int_1^{2^N} \ln x \ln \left( \frac{2^{N+2}}{x} \right) dx + o(\epsilon^{-2}) \\
 &= c_1 I_1 - c_1 I_2 + o(\epsilon^{-2}).
 \end{aligned}$$

Since  $2^N \epsilon^2 / N^2$  is bounded above and below independent of  $N$  and  $\epsilon$ , we can write

$$\begin{aligned}
 (4.6) \quad I_1 &\geq \int_2^{2^N} (\ln x)^2 dx + \ln c_2 \int_2^{2^N} \ln x dx \\
 &\quad - \ln \left( \frac{2^N}{N^2} \right) \int_2^{2^N} \ln x dx - 2 \int_2^{2^N} \ln x \ln \ln x dx.
 \end{aligned}$$

Now  $\int (\ln x)^2 = x(\ln x)^2 - 2x \ln x + 2x$ ;  $\int \ln x = x \ln x - x$ ; and  $\int \ln x \ln \ln x = x \ln x \ln \ln x - x - \int \ln \ln x$ . Carrying out the integrations in (4.6) and using the fact that  $|\int_2^{2^N} \ln \ln x dx| \leq 2^N \ln \ln 2^N = o(N 2^N)$ , we get

$$(4.7) \quad I_1 \geq -cN 2^N.$$

For  $I_2$ , we have

$$\begin{aligned}
 (4.8) \quad I_2 &\leq c 2^N \int_{2^{-N}}^1 \ln(2^N u) \ln(4/u) du \\
 &= c 2^N \ln 2^N \int_{2^{-N}}^1 \ln(4/u) du + c 2^N \int_{2^{-N}}^1 \ln u \ln(4/u) du \\
 &\leq cN 2^N.
 \end{aligned}$$

Combining, we get

$$(4.9) \quad q_1 \geq -cN2^N = -cN^3 2^N / N^2 \geq -c(\ln(1/\epsilon))^3 \epsilon^{-2}.$$

For  $\gamma \geq \gamma_0 > 0$ ,

$$\ln P(|Z| < \gamma) = \ln(1 - 2P(Z > \gamma)) \geq \ln(1 - c_1 \exp(-c_2 \gamma^2)) \geq -c_1 \exp(-c_2 \gamma^2),$$

for constants depending on  $\gamma_0$  but not  $\gamma$ . Hence

$$(4.10) \quad \begin{aligned} q_2 &= \sum_{\{\mathbf{m}: \psi(\mathbf{m}) > N\}} \sum_{j \in \Gamma(\mathbf{m})} r_{j\mathbf{m}} \\ &= \sum_{k=N+1}^{\infty} \sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} 2^k \ln P\left(|Z| < \frac{2^{k/2} \beta_{\mathbf{m}} \epsilon}{2}\right) \\ &\geq -c_1 \sum_{k=N+1}^{\infty} (k+3) 2^k \exp\left(\frac{-c_2 2^k \epsilon^2}{(k+4)^2 (k+1-N)^4}\right) \\ &\geq -c_1 N^2 \epsilon^{-2} \sum_{j=1}^{\infty} (j+N) 2^j \exp\left(\frac{-c_2 2^j N^2}{(j+N)^2 (j+1)^4}\right) \\ &= -c_1 N^2 \epsilon^{-2} \left(\sum_{j=1}^N + \sum_{j>N}\right) = J_1 + J_2. \end{aligned}$$

Now

$$J_1 \geq -c_1 N^2 \epsilon^{-2} (2N) \sum_{j=1}^{\infty} 2^j \exp\left(\frac{-c_2 2^j}{(j+1)^4}\right) \geq -c_1 N^3 \epsilon^{-2},$$

while

$$J_2 \geq -c_1 N^2 \epsilon^{-2} \sum_{j=1}^{\infty} (2j) 2^j \exp\left(\frac{-c_2 2^j}{j^2 (j+1)^4}\right) \geq -c_1 N^2 \epsilon^{-2}.$$

Substituting in (4.10),

$$(4.11) \quad q_2 \geq -cN^3 \epsilon^{-2} \geq -c\epsilon^{-2} (\ln(1/\epsilon))^3.$$

Adding (4.9) and (4.11) and substituting in (4.4) proves the proposition.  $\square$

**REMARK.** Again, for  $d > 2$ , we get a  $(\ln(1/\epsilon))^{3(d-1)}$  term from the fact that  $\sum_{\{\mathbf{m}: \psi(\mathbf{m})=k\}} 1 \approx k^{d-1}$ . To use the two different estimates for  $P(|Z| < \gamma)$  and yet have  $\sum \beta_{\mathbf{m}} \leq 1$ , we must choose  $N$  so that  $2^N / N^{2(d-1)} \approx \epsilon^{-2}$ .

Finally, we have

**THEOREM 4.5.** Suppose  $F$  and  $G$  have densities  $f$  and  $g$ , respectively, with  $d(f^{1/2})/dt \in L^1([0, 1]^d, dt)$  and  $g/\sqrt{f} \in L^2([0, 1]^d, dt)$ . Suppose  $G(\mathbf{1}) = 0$ .

Then

$$P(\|W_F - G\| \leq \varepsilon) \geq c_1 \exp\left(-\frac{1}{2} \int_{[0,1]^d} (g^2/f) dt - c_2 \varepsilon^{-2} (\ln(1/\varepsilon))^{3(d-1)}\right),$$

where  $c_1$  and  $c_2$  are constants depending only on  $F$ .

PROOF. Combine Propositions 4.1, 4.2, 4.3 and 4.4.  $\square$

### REFERENCES

- ADLER, R. J. and BROWN, L. D. (1986). Tail behaviour for suprema of empirical processes. *Ann. Probab.* **14** 1–30.
- BLUM, J. R., KIEFER, J. and ROSENBLATT, M. (1961). Distribution free tests of independence based on the sample distribution function. *Ann. Math. Statist.* **32** 485–492.
- BOROVKOV, A. A. (1967). Boundary value problems for random walks and large deviations in function spaces. *Theory Probab. Appl.* **12** 575–595.
- CABAÑA, E. M. and WSCHBOR, M. (1982). The two-parameter Brownian bridge: Kolmogorov's inequalities and upper and lower bounds for the distribution of the maximum. *Ann. Probab.* **10** 289–301.
- CAIROLI, R. (1970). Une inégalité pour martingales à indices multiples et ses applications. *Séminaire de Probabilités IV. Lecture Notes in Math.* **124** 1–27. Springer, Berlin.
- CSÁKI, E. (1980). A relation between Chung's and Strassen's law of the iterated logarithm. *Z. Wahrsch. verw. Gebiete* **54** 287–301.
- GOODMAN, V. (1976). Distribution estimates for functionals of the two-parameter Wiener process. *Ann. Probab.* **4** 977–982.
- GROENEBOOM, P., OOSTERHOFF, J. and RUYMGAART, F. H. (1979). Large deviation theorems for empirical probability measures. *Ann. Probab.* **7** 553–586.
- PARK, W. J. (1970). A multiparameter Gaussian process. *Ann. Math. Statist.* **41** 1582–1595.
- RÉVÉSZ, P. (1981). How small are the increments of a Wiener sheet. *First Pannonian Symp. Math. Statist. Lecture Notes in Statist.* **8** 207–219. Springer, Berlin.
- SANOV, I. N. (1957). On the probability of large deviations of random variables. *Mat. Sb.* **42** 11–44. (English translation in *Select. Transl. Math. Statist. Probab.* **1** (1961) 213–244.)
- VARADHAN, S. R. S. (1984). *Large Deviations and Applications*. SIAM, Philadelphia.

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF WASHINGTON  
SEATTLE, WASHINGTON 98195