



**Brownian Motion with Lower Class Moving Boundaries Which Grow Faster Than
t^{1/2}**

R. F. Bass; M. Cranston

The Annals of Probability, Vol. 11, No. 1. (Feb., 1983), pp. 34-39.

Stable URL:

<http://links.jstor.org/sici?sici=0091-1798%28198302%2911%3A1%3C34%3ABMWLCM%3E2.0.CO%3B2-3>

The Annals of Probability is currently published by Institute of Mathematical Statistics.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ims.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

BROWNIAN MOTION WITH LOWER CLASS MOVING BOUNDARIES WHICH GROW FASTER THAN $t^{1/2}$

BY R. F. BASS AND M. CRANSTON

University of Illinois, Urbana

Upper and lower bounds are obtained for $P(|W(t)| \leq f(t), t \leq u)$ and $P(|S(n)| \leq f(n), n \leq N)$, u, N large, where $W(t)$ is a Brownian motion, $S(n)$ is a random walk with $ES(1) = 0, E|S(1)|^{2+2\eta} < \infty$, and $f(t)$ is a deterministic function growing faster than $t^{1/2}$ but slower than $(2t \ln \ln t)^{1/2}$.

1. Introduction. Let $W(t)$ be a standard Brownian motion, $f(t)$ a deterministic increasing function. How does $\mathcal{P}_u = P(|W(t)| \leq f(t), t \leq u)$ behave for large u ? A closely related problem is the behavior of $\mathcal{P}_N = P(|S(n)| \leq f(n), n \leq N)$, where $S(n)$ is a random walk.

When $f(t) = ct^{1/2}$, this problem has been considered by Breiman (1965), Gundy and Siegmund (1967), and Brown (1969), among others. When $f(t) = o(t^{1/2})$, results have been obtained by Lai (1977), Portnoy (1978), Kesten (1978), and Novikov (1981), to mention only a few. Upper class functions f have been studied by Cuzick (1981) and Jennen and Lerche (1981).

Here we consider the remaining case, when f grows faster than $t^{1/2}$ but more slowly than $(2t \ln \ln t)^{1/2}$. In contrast to the $o(t^{1/2})$ case, where $\ln \mathcal{P}_u$ is asymptotically $-c \int_0^u f(t)^{-2} dt$, we get that $\ln \mathcal{P}_u$ behaves like $-\int_0^u f(t)^{-2} \exp(-cf(t)^2/2t) dt$ and a similar result for random walks.

More precisely, we get the following:

Suppose $f(t)$ is bounded away from 0, and for $t \geq u_1 > 0$, for some u_1 , $f(t) = t^{1/2}L(t)$, where $L(t)$ is strictly positive, nondecreasing, continuous, slowly varying, $L(t) = o((\ln \ln t)^{1/2})$, and $L(+\infty) = +\infty$. Let $\alpha = 1/3000$.

THEOREM 1. *Given $\varepsilon > 0$, there exist constants c_1, c_2 , and u_0 , depending only on ε and f , such that if $u \geq u_0$,*

$$c_1 \exp\left(-\int_0^u f(t)^{-2} e^{-\alpha f(t)^2/2t} dt\right) \leq \mathcal{P}_u \leq c_2 \exp\left(-\int_0^u f(t)^{-2} e^{-(1+\varepsilon)f(t)^2/2t} dt\right).$$

THEOREM 2. *Let $X_i, i = 1, 2, \dots$ be iid random variables, $S(n) = \sum_{i=1}^n X_i$. Suppose $EX_1 = 0, EX_1^2 = 1$, and $E|X_1|^{2+2\eta} < \infty$ for some $\eta > 0$. Then there exist constants c_1, c_2 , and N_0 such that if $N \geq N_0$,*

$$c_1 \exp\left(-\sum_{i=1}^N f(i)^{-2} e^{-\alpha f(i)^2/2i}\right) \leq \mathcal{P}_N \leq c_2 \exp\left(-\sum_{i=1}^N f(i)^{-2} e^{-(1+\varepsilon)f(i)^2/2i}\right).$$

Our techniques still work for $f(t) = c(2t \ln \ln t)^{1/2}$, but the results are of no interest unless $c < \alpha$.

Theorem 1 is proved in Section 2. Theorem 2 is proved in Section 3.

2. Brownian Motion. We start by getting an estimate on $Q(t, x, k) = P^x(|W(s)| \leq k, s \leq t)$. Note that $Q(t, -x, k) = Q(t, x, k)$ by symmetry.

Let $d_\varepsilon \geq 4$ be such that $d^{-1} \exp(-d^2/2) \leq d^{-2} \exp(-(1-\varepsilon)d^2/2)$ whenever $d \geq d_\varepsilon$.

Received November 1981; revised June 1982.

AMS 1970 Subject Classification. Primary 60J65, 60J15; secondary 60G40.

Key Words and Phrases. Moving boundaries, Brownian motion, random walks.

PROPOSITION 2.1. Suppose $0 \leq x \leq k$. If $(k-x)^2/t \geq d_\varepsilon^2$ then

$$1 - (4t(2\pi)^{-1/2}/(k-x)^2) \exp(-(1-\varepsilon)(k-x)^2/2t) \leq Q(t, x, k) \leq 1 - (t(2\pi)^{-1/2}/k^2) \exp(-k^2/2t).$$

PROOF OF 2.1. Letting $d = k/t^{1/2}$ and using the inequality

$$\int_d^\infty e^{-y^2/2} dy \geq (d^{-1} - d^{-3})e^{-d^2/2} \geq \frac{1}{2} d^{-1} e^{-d^2/2}$$

for $d \geq 2$, (Feller, 1968, page 175) we have

$$Q(t, x, k) \leq P^x(|W(t)| \leq k) \leq P^0(|W(t)| \leq k) = 1 - 2(2\pi t)^{-1/2} \int_k^\infty e^{-y^2/2t} dy \leq 1 - (2\pi)^{-1/2} d^{-1} e^{-d^2/2}$$

if $k^2/t \geq 4$. Since $d^{-1} \geq d^{-2}$ for $d \geq 1$, this gives the right hand inequality.

Let $q(t, x, y, k)$ be the transition density for Brownian motion started at x and killed on leaving $[-k, k]$. Of course, $Q(t, x, k) = \int_{-k}^k q(t, x, y, k) dy$.

A simple change of variables applied to a formula of Feller (1971), page 341, gives

$$(2\pi t)^{1/2} q(t, x, y, k) = \sum_{j=-\infty}^\infty [\exp(-(y-x+4jk)^2/2t) - \exp(-(y+x+4jk+2k)^2/2t)].$$

The infinite series on the right is absolutely convergent and of alternating sign. Pairing terms, if $j \geq 1$, $\exp(-(y-x+4jk)^2/2t) - \exp(-(y+x+4jk+2k)^2/2t) \geq 0$ since $x \geq -k$. If $j \leq -1$, $\exp(-(y-x+4jk)^2/2t) - \exp(-(y+x+4(j-1)k+2k)^2/2t) \geq 0$ since $x \leq k$. Therefore

$$(2\pi t)^{1/2} q(t, x, y, k) \geq e^{-(y-x)^2/2t} - e^{-(y+x-2k)^2/2t} - e^{-(y+x+2k)^2/2t}.$$

Integrating y from $-k$ to k ,

$$\begin{aligned} Q(t, x, k) &\geq P^x(|W(t)| \leq k) - P^{2k-x}(|W(t)| \leq k) - P^{-x-2k}(|W(t)| \leq k) \\ &\geq 1 - P^0(W(t) > k-x) - P^0(W(t) < -k-x) \\ &\quad - P^0(-3k-x \leq W(t) \leq -k-x) - P^0(k-x \leq W(t) \leq 3k-x) \\ &\geq 1 - 4P^0(W(t) \geq k-x). \end{aligned}$$

Let $d = (k-x)/t^{1/2}$ and use the inequality

$$\int_d^\infty e^{-y^2/2} dy \leq d^{-1} e^{-d^2/2}$$

to get $Q(t, x, k) \geq 1 - (4/(2\pi)^{1/2}d)e^{-d^2/2}$. If $d \geq d_\varepsilon$, we thus get the left hand inequality. \square

PROOF OF THEOREM 1.

Upper bound. Let $0 < \varepsilon < 1/2$. Let $q > 2$ be chosen so that $(1 - q^{-1})^{-1} \leq 1 + \varepsilon/4$. Let u_0 be chosen equal to q^I for some I and large enough so that if $s \geq u_0/q^3$, $q^2s \geq t \geq qs$, then

- (i) $[f(t) - f(s)]^2/(t-s) > d_\varepsilon$,
- (ii) $f(s)/f(t) \geq (2q)^{-1}$,
- (iii) $(4(2\pi)^{1/2}q^2)^{-1} \exp(-f(t)^2/2(t-s)) \geq \exp(-(1 + \varepsilon/3)f(t)^2/2(t-s))$, and
- (iv) $f(t)^2/2(t-s) \leq (1 + \varepsilon/3)f(s)^2/2s$.

This can be done since $L(t)$ is slowly varying, but increasing to infinity.

Suppose $u \geq u_0$. Let n be the largest integer such that $q^n \leq u$. Let $t_0 = 0$, $t_i = q^i$ for $1 \leq i < n$, $t_n = u$. Let

$$A_i = \{|W(t)| \leq f(t_i), t_{i-1} < t \leq t_i\}, \quad i = 1, \dots, n,$$

and let

$$r_i = (t_i - t_{i-1}) / ((2\pi)^{1/2} f(t_i)^2 \exp(-f(t_i)^2 / 2(t_i - t_{i-1}))).$$

Note $q^{-2} \leq t_{i-1}/t_i \leq q^{-1}$.

Using the Markov property,

$$\begin{aligned} \mathcal{P}_u &\leq P^0(A_1 A_2 \dots A_n) = E^0(P^{W(t_{n-1})}(|W(t)| \leq f(t_n), 0 < t \leq t_n - t_{n-1}); A_1 \dots A_{n-1}) \\ &= E^0(Q(t_n - t_{n-1}, W(t_{n-1}), f(t_n)); A_1 \dots A_{n-1}) \\ &\leq (1 - r_n) P^0(A_1 \dots A_{n-1}). \end{aligned}$$

To get the last inequality we used (i) and (2.1). Repeating

$$\mathcal{P}_u \leq \prod_{i=I+1}^n (1 - r_i) P^0(A_1 \dots A_I),$$

or

$$\ln \mathcal{P}_u \leq c + \sum_{i=I+1}^n \ln(1 - r_i) \leq c - \sum_{i=I+1}^n r_i.$$

Using (ii), (iii), and (iv), if $t_{i-1} < t \leq t_i$

$$\begin{aligned} -r_i &\leq -(4q^2(t_i - t_{i-1})/f(t_i)^2) \exp(-(1 + \varepsilon/3)f(t_i)^2/2(t_i - t_{i-1})) \\ &\leq -((t_i - t_{i-1})/f(t_{i-1})^2) \exp(-(1 + \varepsilon/3)^2 f(t_{i-1})^2/2(t_i - t_{i-1})). \end{aligned}$$

Thus,

$$\ln \mathcal{P}_u \leq c - \int_{t_I}^u f(t)^{-2} \exp(-(1 + \varepsilon)f(t)^2/2t) dt = c' - \int_0^u f(t)^{-2} \exp(-(1 + \varepsilon)f(t)^2/2t) dt.$$

Lower bound. Let $q = 9$, $\alpha = 1/3000$. Let u_0 be chosen equal to q^I for some I and large enough so that if $s \geq u_0/q^3$, $q^2 s \geq t \geq qs$, then

$$(v) [f(s) - f(s/q)]^2/(t - s) \geq 2\alpha f(t)^2/t,$$

$$(vi) [f(s) - f(s/q)]^2/(t - s) \geq d_{1/4}^2 \quad (> 10),$$

$$(vii) (16q^2)^{-1} \exp(-\frac{1}{2}[f(s) - f(s/q)]^2/2(t - s)) \geq (8/(2\pi)^{1/2}) \exp(-\frac{3}{4}[f(s) - f(s/q)]^2/2(t - s)),$$

and

$$(viii) (16q^2)^{-1} [f(s) - f(s/q)]^{-2} \leq f(t)^{-2}.$$

To show that u_0 can be chosen so that (v) holds, observe that the right hand side of (v) is $2\alpha L(t)^2$. But since $f(s/q)/f(s) = q^{-1/2} L(s/q)/L(s) \leq 2q^{-1/2}$ for s sufficiently large and $t/s - 1 \leq q^2$, the left side of (v) is

$$L(s)^2 [1 - f(s/q)/f(s)]^2 / (t/s - 1) \geq L(s)^2 [1 - 2q^{-1/2}]^2 / q^2 \geq 4\alpha L(s)^2 > 2\alpha L(t)^2$$

for s sufficiently large.

Since $L(t)$ increases to ∞ , it follows easily from (v) that u_0 can be chosen so that (vi) and (vii) hold. The inequality (viii) is argued in a manner similar to (v).

Observe that $-2x \leq \ln(1 - x)$ if $0 \leq x \leq \frac{1}{2}$ and that $xe^{-x/2} \leq \frac{1}{4}$ if $x \geq 7$.

Suppose $u \geq u_0$. Let n be the largest integer such that $q_n \leq u$, let $t_0 = 0$, $t_i = q^i$ for $1 \leq i < n$, $t_n = u$,

$$B_i = \{|W(t)| \leq f(t_{i-1}), t_{i-1} < t \leq t_i\}, \quad i = 1, \dots, n,$$

and let

$$v_i = (4(2\pi)^{-1/2}(t_i - t_{i-1})/[f(t_{i-1}) - f(t_{i-2})]^2) \times \exp(-3/4[f(t_{i-1}) - f(t_{i-2})]^2/2(t_i - t_{i-1})).$$

Note $t_{i-2} = t_{i-1}/q$.

$$\begin{aligned} \mathcal{P}_u &\geq P^0(B_1 B_2 \cdots B_n) \\ &= E^0(P^{W(t_{n-1})}(|W(t)| \leq f(t_{n-1}), 0 < t \leq t_n - t_{n-1}); B_1 \cdots B_{n-1}) \\ &= E^0(Q(t_n - t_{n-1}, W(t_{n-1}), f(t_{n-1})); B_1 \cdots B_{n-1}) \\ &\geq (1 - v_n)P^0(B_1 \cdots B_{n-1}) \end{aligned}$$

by (2.1), (vi), the fact that e^{-x}/x is decreasing, and the fact that on B_{n-1} , $|W(t_{n-1})| \leq f(t_{n-2})$. Repeating, $\mathcal{P}_u \geq \prod_{i=I+1}^n (1 - v_i)P^0(B_1 \cdots B_I)$, or

$$\ln \mathcal{P}_u \geq c + \sum_{i=I+1}^n \ln(1 - v_i) \geq c - 2 \sum_{i=I+1}^n v_i,$$

using (vi) again.

Using (v), (vii), and (viii), if $t_{i-1} < t \leq t_i$,

$$\begin{aligned} -2v_i &\geq -((16q^2)^{-1}(t_i - t_{i-1})/[f(t_{i-1}) - f(t_{i-2})]^2) \\ &\quad \times \exp(-1/2[f(t_{i-1}) - f(t_{i-2})]^2/2(t_i - t_{i-1})) \\ &\geq -((t_i - t_{i-1})/f(t_i)^2) \exp(-\alpha f(t_i)^2/2t_i), \end{aligned}$$

or

$$\ln \mathcal{P}_u \geq c - \int_{t_I}^u f(t)^{-2} \exp(-\alpha f(t)^2/2t) dt = c' - \int_0^u f(t)^{-2} \exp(-\alpha f(t)^2/2t) dt. \quad \square$$

COMMENT. The value of α comes from (v). Replacing the 2's in the derivation of (v) by $1 + \delta$, δ small, and varying q allow one to improve the value of α . But the best value of α one could possibly hope for from our method would be

$$\begin{aligned} \sup_q \limsup_{t=q, s \rightarrow \infty} [1 - f(s/q)/f(s)]^2/(t/s - 1) \\ \leq \sup_q (1 - q^{-1/2})^2/(q - 1) = 1/2(5\sqrt{5} - 11) \approx .09, \end{aligned}$$

which is still substantially less than 1. The difficulty comes from the fact that in the proof of the lower bound, $|W(t_{n-1})|$ is close to $f(t_{n-2})$ with nonnegligible probability.

3. Random walk. Let X_i , $i = 1, 2, \dots$ be iid, mean 0, variance 1 random variables with $E|X_1|^{2+2\eta} < \infty$ for some $\eta > 0$. Let $S(n) = \sum_{i=1}^n X_i$. Define $Q(N, x, k) = P^x(|S(n)| \leq k, n \leq N)$.

PROPOSITION 3.1. Let $1/2 > \varepsilon > 0$, $k \geq |x| \geq 0$. There exist constants M and N_0 such that if $N \geq N_0$, $(k - |x|)/N^{1/2} \geq M$, and $k \leq (2N \ln \ln N)^{1/2}$, then

$$\begin{aligned} 1 - (N/(k - |x|)^2) \exp(-(1 - \varepsilon)(k - |x|)^2/2N) \\ \leq Q(N, x, k) \leq 1 - (N/k^2) \exp(-(1 + \varepsilon)k^2/2N). \end{aligned}$$

PROOF. Suppose $x \geq 0$. The other case is similar.

Use Skorokhod imbedding to find a Brownian motion $W(t)$ and stopping times U_1, \dots, U_N such that $U_1, U_2 - U_1, \dots, U_N - U_{N-1}$ are independent, and $(S(1), \dots, S(N))$ is equal in law to $(W(U_1), \dots, W(U_N))$, and $EU_1^{1+\eta} < \infty$ (see Skorokhod, 1965, for example).

If we let $U_0 = 0$, $Y_i = U_i - U_{i-1} - 1$, $i = 1, \dots, N$, the Y_i 's are iid random variables, $EY_1 = EU_1 - 1 = EW(U_1)^2 - 1 = EX_1^2 - 1 = 0$, and $EY_1^{1+\eta} < \infty$.

Then by Petrov (1975), page 283, for each δ , $P(|\sum_{i=1}^N Y_i|/N > \delta) = o(N^{-\eta})$. So for N sufficiently large, $P^0(|U_N - N| > \delta N) \leq N^{-\eta}$.

$$\begin{aligned} P^x(|S(n)| \leq k, n \leq N) &= P^x(|W(U_n)| \leq k, n \leq N) \leq P^x(|W(U_N)| \leq k) \\ &\leq P^x(|W(U_N)| \leq k, |U_N - N| \\ &\leq \delta N, \sup_{|t-N| \leq \delta N} |W(t) - W(N)| < \varepsilon k/3) \\ &\quad + P^x(\sup_{|t-N| \leq \delta N} |W(t) - W(N)| \\ &\geq \varepsilon k/3) + P^x(|U_N - N| > \delta N) \\ &\leq P^x(|W(N)| < (1 + \varepsilon/3)k) + 2P^0(\sup_{t \leq \delta N} |W(t)| > \varepsilon k/3) \\ &\quad + P^0(|U_N - N| > \delta N). \end{aligned}$$

As in the proof of (2.1), the first term on the right is

$$\begin{aligned} &\leq 1 - (N/((2\pi)^{1/2}(1 + \varepsilon/3)^2 k^2)) \exp(-(1 + \varepsilon/3)^2 k^2/2N) \\ &\leq 1 - (3N/k^2) \exp(-(1 + \varepsilon)k^2/2N), \end{aligned}$$

if k^2/N is large enough. The second term is

$$\begin{aligned} 2(1 - P^0(|W(s)| \leq \varepsilon k/3, s \leq \delta N)) &\leq (72 N/((2\pi)^{1/2} \varepsilon^2 k^2)) \exp(-(1 - \varepsilon/3)\varepsilon^2 k^2/(18\delta N)) \\ &\leq (N/k^2) \exp(-(1 + \varepsilon)k^2/2N) \end{aligned}$$

if we take $\delta \leq \varepsilon^2/72$. Finally, since $k \leq (2N \ln \ln N)^{1/2}$, $(N/k^2) \exp(-(1 + \varepsilon)k^2/2N) \geq \exp(-2k^2/2N) \geq \exp(-2 \ln \ln N) = (\ln N)^{-2}$, which is much larger than $N^{-\eta}$ if N and K^2/N are sufficiently large. So for N, k chosen appropriately, the third term will also be $\leq (N/k^2) \exp(-(1 + \varepsilon)k^2/2N)$.

Summing, $P^x(|S(n)| \leq k, n \leq N) \leq 1 - (N/k^2) \exp(-(1 + \varepsilon)k^2/2N)$. To get the other inequality

$$\begin{aligned} P^x(|W(t)| \leq k, t \leq (1 + \varepsilon/3)N) &\leq P^x(|W(t)| \leq k, t \leq U_N) + P^x(U_N > (1 + \varepsilon/3)N) \\ &\leq P^x(|W(U_n)| \leq k, n \leq N) + P^0(|U_N - N| > \varepsilon N/3), \end{aligned}$$

or

$$\begin{aligned} P^x(|S(n)| \leq k, n \leq N) &= P^x(|W(U_n)| \leq k, n \leq N) \\ &\geq P^x(|W(t)| \leq k, t \leq (1 + \varepsilon/3)N) - P^0(|U_N - N| > \varepsilon N/3). \end{aligned}$$

The first term on the right, by (2.1), is

$$\begin{aligned} &\geq 1 - (4(1 + \varepsilon/3)N/((2\pi)^{1/2}(k-x)^2)) \exp(-(1 + \varepsilon/3)(k-x)^2/2(1 + \varepsilon/3)N) \\ &\geq 1 - (N/2(k-x)^2) \exp(-(1 - \varepsilon)(k-x)^2/2N) \end{aligned}$$

if $(k-x)^2/N$ is large, while the second term is $\leq N^{-\eta}$ for N large, which in turn is $\leq (N/2(k-x)^2) \exp(-(1 - \varepsilon)(k-x)^2/2N)$ by the upper bound on k , as above.

Hence for N large, $(k-x)/N^{1/2}$ large,

$$Q(N, x, k) \geq 1 - (N/(k-x)^2) \exp(-(1 - \varepsilon)(k-x)^2/2N).$$

PROOF OF THEOREM 2. Let $q = 9$ in the lower bound case, q a large integer in the upper bound case. Using (3.1) in place of (2.1), the proof is virtually identical to the proof of Theorem 1. \square

REFERENCES

- [1] BREIMAN, L. (1965). First exit times from a square root boundary. Proc. Fifth Berkeley Symp., University of California Press, Berkeley.

- [2] BROWN, B. M. (1969). Moments of a stopping rule related to the Central Limit Theorem. *Ann. Math. Statist.* **40** 1236–1249.
- [3] FELLER, W. (1971). *An Introduction to Probability Theory and its Applications*, Vol. 2. Wiley, New York.
- [4] GUNDY, R. F. and SIEGMUND, D. (1967). On a stopping rule and the Central Limit Theorem. *Ann. Math. Statist.* **38** 1915–1917.
- [5] KESTEN, H. (1978). Branching Brownian motion with absorption. *Stochastic Processes Appl.* **7** 9–48.
- [6] LAI, T. L. (1977). First exit times from moving boundaries for sums of independent random variables. *Ann. Probability* **5** 210–221.
- [7] NOVIKOV, A. A. (1981). On estimates and the asymptotic behavior of nonexit probabilities of a Wiener process to a moving boundary. *Math. USSR Sbornik* **38** 495–505.
- [8] PETROV, V. V. (1975). *Sums of Independent Random Variables*. Springer-Verlag, New York.
- [9] PORTNOY, S. (1978). Probability bounds for first exits through moving boundaries, *Ann. Probability* **6** 106–117.
- [10] SKOROKHOD, A. V. (1965). *Studies in the Theory of Random Processes*. Addison-Wesley, Reading, Massachusetts.
- [11] FELLER, W. (1968). *An Introduction to Probability Theory and its Applications*, Vol. 1. Wiley, New York.
- [12] CUZICK, J. (1981). Boundary crossing probabilities for stationary Gaussian processes and Brownian motion. *Trans. Amer. Math. Soc.* **263** 469–492.
- [13] JENNEN, C. and LERCHE, H. R. (1981). First exit densities of Brownian motion through one-sided moving boundaries. *Z. Wahrsch. verw Gebiete* **55** 133–148.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF WASHINGTON
SEATTLE, WA 98195

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ROCHESTER
ROCHESTER, NY 14627