

OPERATOR ALGEBRAS AND MAULDIN-WILLIAMS GRAPHS

MARIUS IONESCU

ABSTRACT. We describe a method for associating a C^* -correspondence to a Mauldin-Williams graph and show that the Cuntz-Pimsner algebra of this C^* -correspondence is isomorphic to the C^* -algebra of the underlying graph. In addition, we analyze certain ideals of these C^* -algebras.

We also investigate Mauldin-Williams graphs and fractal C^* -algebras in the context of the Rieffel metric. This generalizes the work of Pinzari, Watatani and Yonetani. Our main result here is a “no go” theorem showing that such algebras must come from the commutative setting.

1. Introduction. In recent years many classes of C^* -algebras have been shown to fit into the Pimsner construction of what are known now as Cuntz-Pimsner algebras, see [20, 22]. This construction is based on a so-called C^* -correspondence over a C^* -algebra. For example, a natural C^* -correspondence can be associated with a graph G , see [10], [11, Example 1.5]. The Cuntz-Pimsner algebra of this C^* -correspondence is isomorphic to the graph C^* -algebra $C^*(G)$ as defined in [16]. Another example is the C^* -correspondence associated with a local homeomorphism on a compact metric space studied by Deaconu in [6], and the C^* -correspondence associated with a local homeomorphism on a locally compact space studied by Deaconu, Kumjian, and Muhly in [7]. They showed that the Cuntz-Pimsner algebra is isomorphic to the groupoid C^* -algebra associated with a local homeomorphism in [5, 7, 26].

By a (directed) *graph* we mean a system $G = (V, E, r, s)$ where V and E are finite sets, called the sets of *vertices* and *edges*, respectively, of the graph, and where r and s are maps from E to V , called the *range* and *source* maps, respectively. Thus, $s(e)$ is the source of an edge e and $r(e)$ is its range. A *Mauldin-Williams graph* is a graph G together with a collection of compact metric spaces, one for each

AMS *Mathematics subject classification.* Primary 26A18, 37A55, 37B10, 37E25, 46L08, 46L55, 46L89.

Received by the editors on Jan. 28, 2004, and in revised form on Nov. 29, 2004.

vertex of the graph, and a collection of contraction maps, one for each edge of the graph which satisfy certain properties, see Definition 2.1 below. In this note we follow the notations from [8]. We associate with such a system a C^* -correspondence which reflects the dynamics of the Mauldin-Williams graph, and we analyze the Cuntz-Pimsner algebra of this C^* -correspondence. Our construction is related with topological generalizations of graph C^* -algebras of Katsura [14] and Muhly and Tomforde [21]. The study of the Cuntz-Pimsner algebra associated with graph dynamical systems was initiated in [23], where the authors consider the case when the graph G consists of a single vertex v and edges e_1, e_2, \dots, e_n . In this case, the ϕ_e s constitute what is known as an *iterated function system* acting on the space (T_v, ρ_v) . They conclude that $\mathcal{O}(\mathcal{X})$ is isomorphic to the Cuntz algebra \mathcal{O}_n . The first of our two principal theorems in this note generalizes this result. Our proof is different from the proof in [23] and reveals extra structure.

Theorem. *Let $(G, \{T_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$ be a Mauldin-Williams graph such that the graph G has no sinks and no sources. Let A and \mathcal{X} be the associated C^* -algebra and C^* -correspondence. Then the Cuntz-Pimsner algebra $\mathcal{O}(\mathcal{X})$ is isomorphic to $C^*(G)$ of [4].*

Thus the structure of $\mathcal{O}(\mathcal{X})$ is completely determined by the graph G . From one perspective, this result is somewhat disappointing. Given the richness of dynamical systems expressed as Mauldin-Williams graphs and given the fact that Cuntz-Pimsner algebras generalize crossed products, one might expect a lively interplay between the dynamics and the structure of $\mathcal{O}(\mathcal{X})$. However, the “rigidity” that this theorem expresses is quite remarkable, and it may inspire one to wonder about the natural limits of the result.

In particular, one might wonder if there are noncommutative versions of Mauldin-Williams graphs and whether these might prove to have a richer theory. This thought was taken up in [23, Section 4.3] where Pinzari, Watatani and Yonetani considered noncommutative iterated function systems based on Rieffel’s notion of “noncommutative metric spaces” [27, 28]. The second objective of this note is to show that noncommutative iterated function systems of Pinzari, Watatani and Yonetani can be formulated in the setting of Mauldin-Williams-type graphs, but that the generality gained is illusory. Roughly, the Rieffel

metric is a metric on the state space of a (not necessarily commutative) C^* -algebra A defined by a certain subset of “Lipschitz elements” in A (see Definition 3.1 below). When we associate a C^* -algebra A_v to each vertex $v \in V$, for a prescribed graph $G = (V, E, r, s)$, and when we associate a $*$ -homomorphism $\phi_e : A_{s(e)} \rightarrow A_{r(e)}$ to each edge $e \in E$ that is strictly contractive with respect to the Rieffel metrics on $A_{s(e)}$ and $A_{r(e)}$, we call the resulting system a noncommutative Mauldin-Williams graph. It also gives rise to a Cuntz-Pimsner algebra $\mathcal{O}(\mathcal{X})$. Our second objective in this note is to show that once more $\mathcal{O}(\mathcal{X})$ is isomorphic to the Cuntz-Krieger algebra associated to G . In fact, we shall show in Theorem 3.4 that, in such situations, the C^* -algebras A_v are necessarily commutative. This implies, in particular, that the structures considered by Pinzari, Watatani and Yonetani are necessarily no more general than those arising from ordinary iterated function systems.

2. The Cuntz-Pimsner algebra associated to a Mauldin-Williams graph.

Definition 2.1. By a *Mauldin-Williams graph*, see [8, 18], we mean a system $(G, \{T_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$ where $G = (V, E, r, s)$ is a graph and where $\{T_v, \rho_v\}_{v \in V}$ and $\{\phi_e\}_{e \in E}$ are families such that:

- (1) For each $v \in V$, T_v is a compact metric space with a prescribed metric ρ_v .
- (2) For $e \in E$, ϕ_e is a continuous map from $T_{r(e)}$ to $T_{s(e)}$ such that

$$\rho_{s(e)}(\phi_e(x), \phi_e(y)) \leq c\rho_{r(e)}(x, y)$$

for some constant c satisfying $0 < c < 1$, independent of e , and all $x, y \in T_{r(e)}$.

We shall assume, too, that the functions s and r are surjective. Thus, we assume that there are no sinks and no sources in the graph G . An *invariant list* associated with a Mauldin-Williams graph is a family $(K_v)_{v \in V}$ of compact sets, such that $K_v \subset T_v$ for all $v \in V$ and

$$K_v = \bigcup_{\substack{e \in E \\ s(e)=v}} \phi_e(K_{r(e)}).$$

Since each ϕ_e is a contraction, a Mauldin-Williams graph

$$(G, \{T_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$$

has a unique invariant list, see [18, Theorem 1]. We set $T := \cup_{v \in V} T_v$ and $K := \cup_{v \in V} K_v$, and we call K the *invariant set* of the Mauldin-Williams graph.

In the particular case when we have one vertex v and n edges, i.e., in the setting of an *iterated function system*, the invariant set is the unique compact subset $K := K_v$ of $T = T_v$ such that

$$K = \phi_1(K) \cup \dots \cup \phi_n(K).$$

Definition 2.2. Given a Mauldin-Williams graph

$$(G, \{T_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E}),$$

we construct a so called C^* -correspondence \mathcal{X} over the C^* -algebra $A = C(T)$, where $T = \coprod_{v \in V} T_v$ is the disjoint union of the spaces T_v , as follows. Let $E \times_G T = \{(e, x) : | : x \in T_{r(e)}\}$. Then, by our finiteness assumptions, $E \times_G T$ is a compact space. We set $\mathcal{X} = C(E \times_G T)$ and view \mathcal{X} as a bimodule over $C(T)$ via the formulae:

$$\xi \cdot a(e, x) := \xi(e, x)a(x)$$

and

$$a \cdot \xi(e, x) := a \circ \phi_e(x)\xi(e, x),$$

where $a \in C(T)$ and $\xi \in C(E \times_G T)$. Further, \mathcal{X} comes equipped with the structure of a Hilbert C^* -module over $C(T)$ via the formula

$$\langle \xi, \eta \rangle_A(x) := \sum_{\substack{e \in E \\ x \in T_{r(e)}}} \overline{\xi(e, x)} \eta(e, x)$$

for all $\xi, \eta \in \mathcal{X}$ so that, in the language of [20], \mathcal{X} may be viewed as a C^* -correspondence over $C(T)$. Since there are no sources in the graph

G , the A -valued inner product is well defined. Let $n = |E|$, and let $C^n(A)$ be the column space over A , i.e., $C^n(A) = \{(\xi_e)_{e \in E} : \xi_e \in A, \text{ for all } e \in E\}$. Then we view \mathcal{X} as a subset of $C^n(A)$.

We note that the left action is given by the $*$ -homomorphism $\Phi : A \rightarrow \mathcal{L}(\mathcal{X})$, $(\Phi(a)\xi)(e, x) = a \circ \phi_e(x)\xi(e, x)$. Then Φ is faithful if and only if $K = T$.

In the case of an iterated function system the C^* -correspondence will be the full column space over the C^* -algebra $A = C(T)$, that is $\mathcal{X} = C^n(A)$, with the structure:

- The right action is the untwisted right multiplication, i.e., $\xi \cdot a(i, x) = \xi(i, x)a(x)$ for all $i \in \{1, \dots, n\}$ and $x \in T$.
- The left action is given by the $*$ -homomorphism $\Phi : A \rightarrow \mathcal{L}(\mathcal{X})$ defined by the formula $\Phi(a)(\xi)(i, x) = a \circ \varphi_i(x)\xi_i(x)$.
- The A -valued inner product given by the formula:

$$\langle \xi, \eta \rangle_A(x) = \sum_{i=1}^n \xi^*(i, x) \eta(i, x).$$

Given a C^* -correspondence \mathcal{X} over a C^* -algebra A a *Toeplitz representation* of \mathcal{X} in a C^* -algebra B consists of a pair (ψ, π) , where $\psi : \mathcal{X} \rightarrow B$ is a linear map and $\pi : A \rightarrow B$ is a $*$ -homomorphism such that

$$\psi(x \cdot a) = \psi(x)\pi(a), \quad \psi(a \cdot x) = \pi(a)\psi(x),$$

i.e., the pair (ψ, π) is a bimodule map and

$$\psi(x)^* \psi(y) = \pi(\langle x, y \rangle_A).$$

That is, the map ψ preserves inner product, see [11, Section 1]. Given such a Toeplitz representation, there is an $*$ -homomorphism $\pi^{(1)}$ from $\mathcal{K}(\mathcal{X})$ into B which satisfies

$$\pi^{(1)}(\Theta_{x,y}) = \psi(x)\psi(y)^* \quad \text{for all } x, y \in \mathcal{X},$$

where $\Theta_{x,y} = x \otimes \tilde{y}$ is the rank one operator defined by $\Theta_{x,y}(z) = x \cdot \langle y, z \rangle_A$.

We define then

$$J(\mathcal{X}) := \Phi^{-1}(\mathcal{K}(\mathcal{X})),$$

which is a closed two sided-ideal in A , see [11, Definition 1.1]. Let K be an ideal in $J(\mathcal{X})$. We say that a Toeplitz representation (ψ, π) of \mathcal{X} is *coisometric* on K if

$$\pi^{(1)}(\Phi(a)) = \pi(a) \quad \text{for all } a \in K.$$

When (ψ, π) is coisometric on all of $J(\mathcal{X})$, we say that it is *Cuntz-Pimsner covariant*.

It is shown in [11, Proposition 1.3] that, for an ideal K in $J(\mathcal{X})$, there is a C^* -algebra $\mathcal{O}(K, \mathcal{X})$ and a Toeplitz representation $(k_{\mathcal{X}}, k_A)$ of \mathcal{X} into $\mathcal{O}(K, \mathcal{X})$ which is coisometric on K and satisfies:

(1) for every Toeplitz representation (ψ, π) of \mathcal{X} which is coisometric on K , there is an $*$ -homomorphism $\psi \times_K \pi$ of $\mathcal{O}(K, \mathcal{X})$ such that $(\psi \times_K \pi) \circ k_{\mathcal{X}} = \psi$ and $(\psi \times_K \pi) \circ k_A = \pi$; and

(2) $\mathcal{O}(K, \mathcal{X})$ is generated as a C^* -algebra by $k_{\mathcal{X}}(\mathcal{X}) \cup k_A(A)$.

The algebra $\mathcal{O}(\{0\}, \mathcal{X})$ is the *Toeplitz algebra* $\mathcal{T}_{\mathcal{X}}$, and $\mathcal{O}(J(\mathcal{X}), \mathcal{X})$ is the *Cuntz-Pimsner algebra* $\mathcal{O}_{\mathcal{X}}$.

For a finite graph $G = (E, V, r, s)$, a *Cuntz-Krieger G -family* consists of a family $\{P_v : v \in V\}$ of mutually orthogonal projections and a family of partial isometries $\{S_e\}_{e \in E}$ such that

$$S_e^* S_e = P_{r(e)} \quad \text{for } e \in E, \quad \text{and} \quad P_v = \sum_{s(f)=v} S_f S_f^* \quad \text{for } v \in s(E).$$

The edge matrix of G is the $E \times E$ matrix defined by

$$A_G(e, f) = \begin{cases} 1 & \text{if } r(e) = s(f) \\ 0 & \text{otherwise.} \end{cases}$$

Then, a Cuntz-Krieger G -family satisfies:

$$S_e^* S_e = \sum_{f \in E} A_G(e, f) S_f S_f^*$$

for every $e \in E$ such that $A_G(e, \cdot)$ has nonzero entries. It is shown in [16, Theorem 1.2] that there exists a C^* -algebra $C^*(G)$ generated

by a Cuntz-Krieger G -family $\{S_e, P_v\}$ of nonzero elements such that, for every Cuntz-Krieger G -family $\{W_e, T_v\}$ of partial isometries on H , there is a representation π of $C^*(G)$ on H such that $\pi(S_e) = W_e$ and $\pi(P_v) = T_v$ for all $e \in E$ and $v \in V$. The triple $(C^*(G), S_e, P_v)$ is unique up to isomorphism. Since we are assuming that G has no sinks, $\{S_e\}_{e \in E}$ is a Cuntz-Krieger family for the edge matrix A_G in the sense of [4], see [16, Section 1], and the projections P_v are redundant. If the matrix A_G satisfies Condition (I) from [4] (or, equivalently, since G is finite, if G satisfies Condition (L) from [16], which asserts that every loop has an exit), then $C^*(G)$ is unique and is isomorphic to the Cuntz-Krieger algebra from [4], see [16, Theorem 3.7].

Theorem 2.3. *Let $(G, \{T_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$ be a Mauldin-Williams graph such that the graph G has no sinks and no sources. Let A and \mathcal{X} be defined as above. Then the Cuntz-Pimsner algebra $\mathcal{O}_{\mathcal{X}}$ is isomorphic to $C^*(G)$.*

Before proving the theorem, we introduce some notation. For $k \geq 2$, set

$$E^k := \{\alpha = (\alpha_1, \dots, \alpha_k) : \alpha_i \in E \text{ and } r(\alpha_i) = s(\alpha_{i+1}), i = 1, \dots, k - 1\},$$

the set of *paths of length k* in the graph G . Let $E^* = \cup_{k \in \mathbf{N}} E^k$ be the space of *finite paths* in the graph G . Also the *infinite path space* E^∞ is defined to be

$$E^\infty := \{(\alpha_i)_{i \in \mathbf{N}} : \alpha_i \in E \text{ and } r(\alpha_i) = s(\alpha_{i+1}) \forall i \in \mathbf{N}\}.$$

For $v \in V$, we also define $E^k(v) := \{\alpha \in E^k : s(\alpha) = v\}$, and we define $E^*(v)$ and $E^\infty(v)$ in a similar way. We consider $E^\infty(v)$ endowed with the metric $\delta_v(\alpha, \beta) = c^{|\alpha \wedge \beta|}$ if $\alpha \neq \beta$ and 0 otherwise, where $\alpha \wedge \beta$ is the longest common prefix of α and β , and $|w|$ is the length of the word $w \in E^*$, see [8, p. 116]. Then $E^\infty(v)$ is a compact metric space and, since E^∞ equals the disjoint union of the spaces $E^\infty(v)$, E^∞ becomes a compact metric space in a natural way.

For $\alpha \in E^k$, we write $\phi_\alpha = \phi_{\alpha_1} \circ \dots \circ \phi_{\alpha_k}$ and $S_\alpha = S_{\alpha_1} \dots S_{\alpha_k}$. Let \mathcal{S}_v be the state space of $A_v = C(T_v)$ and $\mathcal{S} = \prod_{v \in V} \mathcal{S}_v$. We consider

the metrics L_v defined on \mathcal{S}_v by the formula

$$(2.1) \quad L_v(\mu, \nu) = \sup\{|\mu(f) - \nu(f)| : f \in \text{Lip}(T_v), c_f \leq 1\},$$

where $\text{Lip}(T_v)$ is the space of Lipschitz functions on T_v and c_f is the Lipschitz constant of the Lipschitz function f . For $f \in \text{Lip}(T_v)$, $v \in V$ and $\mu, \nu \in \mathcal{S}_v$,

$$(2.2) \quad |\mu(f) - \nu(f)| \leq c_f L_v(\mu, \nu).$$

Further, if $\alpha \in E^k$, $k \geq 1$, and $\mu, \nu \in \mathcal{S}_{r(\alpha)}$, then $\mu \circ \phi_{\alpha_k}^{-1} \circ \dots \circ \phi_{\alpha_1}^{-1}, \nu \circ \phi_{\alpha_k}^{-1} \circ \dots \circ \phi_{\alpha_1}^{-1} \in \mathcal{S}_{s(\alpha)}$ and

$$(2.3) \quad L_{s(\alpha_1)}(\mu \circ \phi_{\alpha_k}^{-1} \circ \dots \circ \phi_{\alpha_1}^{-1}, \nu \circ \phi_{\alpha_k}^{-1} \circ \dots \circ \phi_{\alpha_1}^{-1}) \leq c^k \text{diam}_L(\mathcal{S}),$$

where $\text{diam}_L(\mathcal{S}) = \max_{v \in V} \text{diam}_{L_v}(\mathcal{S}_v)$.

For $\alpha \in E^\infty$, the sequence $(\phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)}))_{k \in \mathbf{N}} \subset T_{s(\alpha)}$ is a decreasing sequence of compact sets. Moreover, $\text{diam}(\phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)})) \leq c^k D$, where $D := \max_{v \in V} \text{diam}(T_v)$. Therefore

$$\lim_{k \rightarrow \infty} \text{diam}(\phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)})) = 0,$$

so the intersection $\bigcap_{k \in \mathbf{N}} \phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)})$ consists of a single point, $x_\alpha \in T_{s(\alpha)}$. Hence, we can define a map $\Pi : E^\infty \rightarrow T$ by the formula

$$\Pi(\alpha) = x_\alpha.$$

Then Π is a continuous map and its image is the invariant set K of the Mauldin-Williams graph.

Proof of Theorem 2.3. Let $\mu_0 = (\mu_v^0)_{v \in V} \in \mathcal{S}$ be fixed and $a = \sum_{v \in V}^\oplus a_v \in \text{Lip}(T)$. We define

$$i_A(a) = \lim_{k \rightarrow \infty} \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^*.$$

We prove that i_A is a norm decreasing $*$ -homomorphism from the $*$ -algebra $\text{Lip}(T)$ into $C^*(G)$. Then, since $\text{Lip}(T)$ is a dense $*$ -subalgebra of $A = C(T)$, we can extend i_A to A .

We show first that the limit from the definition of $i_A(a)$ exists. Let $a \in \text{Lip}(T)$, and let $\varepsilon > 0$. Choose $k_0 \in \mathbf{N}$ such that $c^k \text{diam}_L(\mathcal{S})c_a < \varepsilon$ for all $k \geq k_0$. Set $a_k := \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\beta S_\beta^* S_\alpha^*$. Let $k, m \geq k_0$, and suppose that $k > m$. Then

$$\begin{aligned} a_m - a_k &= \sum_{\alpha \in E^m} \sum_{\substack{\beta \in E^{k-m} \\ s(\beta)=r(\alpha)}} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\beta S_\beta^* S_\alpha^* \\ &\quad - \sum_{\alpha \in E^m} \sum_{\substack{\beta \in E^{k-m} \\ s(\beta)=r(\alpha)}} \mu_{r(\beta)}^0(a_{s(\alpha)} \circ \phi_{\alpha\beta}) S_\alpha S_\beta S_\beta^* S_\alpha^* \\ &= \sum_{\alpha \in E^m} \sum_{\substack{\beta \in E^{k-m} \\ s(\beta)=r(\alpha)}} (\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) \\ &\quad - \mu_{r(\beta)}^0(a_{s(\alpha)} \circ \phi_{\alpha\beta})) S_\alpha S_\beta S_\beta^* S_\alpha^* \\ &= \sum_{\alpha \in E^m} \sum_{\substack{\beta \in E^{k-m} \\ s(\beta)=r(\alpha)}} (\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) \\ &\quad - \mu_{r(\beta)}^0 \circ \phi_\beta^{-1}(a_{s(\alpha)} \circ \phi_\alpha)) S_\alpha S_\beta S_\beta^* S_\alpha^*. \end{aligned}$$

Since $|\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) - \mu_{r(\beta)}^0 \circ \phi_\beta^{-1}(a_{s(\alpha)} \circ \phi_\alpha)| < \varepsilon$, by equations (2.2) and (2.3), for all $\alpha \in E^m, \beta \in E^{k-m}$ such that $s(\beta) = r(\alpha)$, and since $S_\alpha S_\beta S_\beta^* S_\alpha^*$ are orthogonal projections, $\|a_m - a_k\| < \varepsilon$, for all $m, k \geq k_0$. So $(a_k)_{k \in \mathbf{N}}$ is a Cauchy sequence, hence convergent. Since $\|a_m\| = \max_{\alpha \in E^m} |\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha)| \leq \|a\|$ for all $m \in \mathbf{N}$, $\|i_A(a)\| \leq \|a\|$ for all $a \in A$.

Next we prove that i_A is a homomorphism. Let $a, b \in \text{Lip}(T)$. Then for each $\alpha \in E^\infty$ there is a point $x_\alpha \in K$ such that $\bigcap_{k \in \mathbf{N}} \phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)}) = \{x_\alpha\}$. Then $\lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}^0(a_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) = a(x_\alpha)$, $\lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}^0(b_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) = b(x_\alpha)$ and $\lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}^0((ab)_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) = a(x_\alpha)b(x_\alpha)$. Let $\varepsilon > 0$. Since $\text{diam}(\phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)})) \leq c^k D$ for all $\alpha \in E^\infty$ and $k \in \mathbf{N}$, there exists some $N \in \mathbf{N}$ such that $|\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) - a(x_\alpha)| < \varepsilon$, $|\mu_{r(\alpha)}^0(b_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) - b(x_\alpha)| < \varepsilon$ and $|\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k} b_{s(\alpha)} \circ \phi_{\alpha_1 \dots \alpha_k}) - a(x_\alpha)b(x_\alpha)| < \varepsilon$ for all $k \geq N$

and for all $\alpha \in E^\infty$. Then

$$\begin{aligned} & \left\| \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha b_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^* \right. \\ & \quad \left. - \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) \mu_{r(\alpha)}^0(b_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^* \right\| \\ & \leq \max_{\alpha \in E^k} |\mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha b_{s(\alpha)} \circ \phi_\alpha) - \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) \mu_{r(\alpha)}^0(b_{s(\alpha)} \circ \phi_\alpha)| \\ & < \varepsilon(1 + \|a\| + \|b\|) \end{aligned}$$

for all $k \geq N$. Thus, $i_A(ab) = i_A(a)i_A(b)$. Hence, i_A is a homomorphism and one can easily see that it is an $*$ -homomorphism.

Further, i_A satisfies the equation

$$(2.4) \quad i_A(a)S_e = S_e i_A(a_{s(e)} \circ \phi_e) \quad \text{for all } a \in A, \text{ and } e \in E,$$

where we extend the map $a_{s(e)} \circ \phi_e$ to all T by setting it to be 0 when $x \notin T_{r(e)}$, since, for $a \in \text{Lip}(T)$, we have:

$$\begin{aligned} i_A(a)S_e &= \left(\lim_{k \rightarrow \infty} \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^* \right) S_e \\ &= \lim_{k \rightarrow \infty} \sum_{\alpha \in E^k, \alpha_1=e} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_e S_{\alpha_2} \cdots S_{\alpha_k} S_{\alpha_k}^* \cdots S_{\alpha_2}^* \\ &= S_e \lim_{k \rightarrow \infty} \sum_{\alpha' \in E^{k-1}(r(e))} \mu_{r(\alpha')}^0(a_{s(e)} \circ \phi_e \circ \phi_{\alpha'}) S_{\alpha'} S_{\alpha'}^* \\ &= S_e i_A(a_{s(e)} \circ \phi_e). \end{aligned}$$

We also define the linear map $i_{\mathcal{X}} : \mathcal{X} \rightarrow C^*(G)$ by the formula

$$i_{\mathcal{X}}(\xi) = \sum_{e \in E} S_e i_A(\xi_e),$$

where $\xi_e \in C(T)$ is defined by $\xi_e(x) = \xi(e, x)$ if $x \in T_{r(e)}$ and 0 otherwise. We have

$$i_{\mathcal{X}}(\xi \cdot a) = \sum_{e \in E} S_e i_A(\xi_e a) = \sum_{e \in E} S_e i_A(\xi_e) i_A(a) = i_{\mathcal{X}}(\xi) i_A(a),$$

$$\begin{aligned} i_{\mathcal{X}}(a \cdot \xi) &= \sum_{e \in E} S_e i_A(a_{s(e)} \circ \phi_e \xi_e) = \sum_{e \in E} S_e i_A(a_{s(e)} \circ \phi_e) i_A(\xi_e) \\ &= \sum_{e \in E} i_A(a) S_e i_A(\xi_e) = i_A(a) i_{\mathcal{X}}(\xi) \end{aligned}$$

and

$$\begin{aligned} i_{\mathcal{X}}(\xi)^* i_{\mathcal{X}}(\eta) &= \left(\sum_{e \in E} S_e i_A(\xi_e) \right)^* \left(\sum_{f \in E} S_f i_A(\eta_f) \right) \\ &= \sum_{e \in E} i_A(\xi_e)^* i_A(\eta_e) = i_A \left(\sum_{e \in E} \xi_e^* \eta_e \right) = i_A(\langle \xi, \eta \rangle_A). \end{aligned}$$

Hence $(i_A, i_{\mathcal{X}})$ is a Toeplitz representation.

Let $J(\mathcal{X}) := \Phi^{-1}(\mathcal{K}(\mathcal{X}))$. Note that $J(\mathcal{X}) = A$ since for $a \in A$ we have

$$\Phi(a)\xi = \sum_{e \in E} \Theta_{x^e, \delta^e}(\xi),$$

where $x^e \in \mathcal{X}$ is defined by $x_f^e = a_{s(e)} \circ \phi_e \delta_f^e$,

$$\delta^e(f, x) = \begin{cases} 1 & \text{if } f = e \text{ and } x \in T_{r(e)} \\ 0 & \text{otherwise.} \end{cases}$$

Then, for $a \in A$, we have

$$\begin{aligned} i_A^{(1)}(\Phi(a)) &= i_A^{(1)} \left(\sum_{e \in E} \Theta_{x^e, \delta^e} \right) = \sum_{e \in E} i_A^{(1)}(\Theta_{x^e, \delta^e}) \\ &= \sum_{e \in E} i_{\mathcal{X}}(x^e) i_{\mathcal{X}}(\delta^e)^* \\ &= \sum_{e \in E} \left(\sum_{f \in E} S_f i_A(x_f^e) \right) \left(\sum_{g \in E} S_g i_A(\delta_g^e) \right)^* \\ &= \sum_{e \in E} (S_e i_A(a_{s(e)} \circ \phi_e)) (i_A(1_{T_{r(e)}}) S_e^*) \\ &= i_A(a) \sum_{e \in E} S_e S_e^* = i_A(a). \end{aligned}$$

Therefore, $(i_A, i_{\mathcal{X}})$ is a Cuntz-Pimsner covariant representation.

For δ^e defined as above, we notice that

$$i_{\mathcal{X}}(\delta^e) = \sum_{f \in E} S_f i_A(\delta_f^e) = S_e i_A(1_{T_{r(e)}}) = S_e.$$

Then $i_{\mathcal{X}}(\mathcal{X}) \cup i_A(A)$ generates $C^*(G)$.

Since $(i_A, i_{\mathcal{X}})$ is a Cuntz-Pimsner covariant representation, there exists a homomorphism $i_{\mathcal{X}} \times i_A$ of $\mathcal{O}_{\mathcal{X}}$ onto $C^*(G)$ such that $(i_{\mathcal{X}} \times i_A) \circ k_{\mathcal{X}} = i_{\mathcal{X}}$ and $(i_{\mathcal{X}} \times i_A) \circ k_A = i_A$. We prove that $i_{\mathcal{X}} \times i_A$ is also injective. Let $\gamma : \mathbf{T} \rightarrow \text{Aut}(\mathcal{O}_{\mathcal{X}})$ defined by $\gamma_z(k_{\mathcal{X}}(\xi)) = z k_{\mathcal{X}}(\xi)$, and let $\gamma_z(k_A(a)) = k_A(a)$ be the gauge action on $\mathcal{O}_{\mathcal{X}}$. Let $\beta : \mathbf{T} \rightarrow \text{Aut}(C^*(G))$ defined by $\beta_z(S_e) = z S_e$ for all $e \in E$ be the gauge action on $C^*(G)$. Therefore, by the definition of i_A and $i_{\mathcal{X}}$, $\beta_z(i_{\mathcal{X}}(\xi)) = z i_{\mathcal{X}}(\xi)$ and $\beta_z(i_A(a)) = i_A(a)$ for all $\xi \in \mathcal{X}$ and $a \in A$. Hence $\beta_z \circ (i_{\mathcal{X}} \times i_A) = (i_{\mathcal{X}} \times i_A) \circ \gamma_z$ for all $z \in \mathbf{T}$. Then the gauge-invariant uniqueness theorem [11, Theorem 4.1] implies that $i_{\mathcal{X}} \times i_A$ is injective. Thus, $C^*(G)$ is isomorphic to the Cuntz-Pimsner algebra associated to the C^* -correspondence \mathcal{X} . \square

Corollary 2.4. *The Cuntz-Pimsner algebra $\mathcal{O}_{\mathcal{X}} = \mathcal{O}(J(\mathcal{X}), \mathcal{X})$ of the C^* -correspondence associated with an iterated function system $(\phi_1, \phi_2, \dots, \phi_n)$ is isomorphic to the Cuntz algebra \mathcal{O}_n .*

If K (the invariant set of the Mauldin-Williams graph) is a proper subset of T , then $U := T \setminus K$ is a nonempty open set of T . Let $I_U := C_0(U)$ be the corresponding ideal in A . Then

$$\mathcal{X}_{I_U} := \{\xi \in \mathcal{X} : \langle \xi, \eta \rangle_A \in I_U \text{ for all } \eta \in \mathcal{X}\}$$

is a right Hilbert I_U -module, and we know that $\mathcal{X}_{I_U} = \mathcal{X}I_U := \{\xi \cdot i : \xi \in \mathcal{X}, i \in I_U\}$, see [11, Section 2]. It follows that $\mathcal{X}_{I_U} = \{\xi \in \mathcal{X} : \xi_e \in C_0(U)\}$ ($\xi_e \in C_0(U)$ means that $\xi(e, x) = 0$ if $x \in K$). We claim that I_U is an \mathcal{X} -invariant ideal in A , i.e., $\Phi(I_U)\mathcal{X} \subset \mathcal{X}I_U$. For $i \in I_U$ and $\xi \in \mathcal{X}$, we have $(\Phi(i)\xi)_e = i \circ \phi_e \xi_e$, and, since $i \in I_U$ and $\phi_e(K_{r(e)}) \subset K_{s(e)}$, $i \circ \phi_e \in I_U$. Hence, $(\Phi(i)\xi)_e \in I_U$. Therefore, I_U is an \mathcal{X} -invariant ideal in A and $\mathcal{X}/\mathcal{X}I_U$ is a C^* -correspondence over $A/I_U \simeq C(K)$, see [11, Lemma 2.3]. Moreover $\mathcal{X}/\mathcal{X}I_U \simeq \mathcal{X}(K)$, where $\mathcal{X}(K) = C(E \times_G K)$ is the C^* -correspondence defined as in Definition 2.2 for the C^* -algebra

$C(K)$. Then the ideal $\mathcal{I}(I_U)$ of $\mathcal{O}_{\mathcal{X}}$ generated by $i_A(I_U)$ is Morita equivalent to $\mathcal{O}_{\mathcal{X}I_U}$, and since $\Phi(A) \subset \mathcal{K}(\mathcal{X})$, $\mathcal{O}_{\mathcal{X}}/\mathcal{I}(I_U) \cong \mathcal{O}_{\mathcal{X}/\mathcal{X}I_U}$, see [11, Corollary 3.3].

Proposition 2.5. *The ideal $\mathcal{I}(I_U)$ generated by $i_A(I_U)$ is equal to 0.*

Proof. Let $a \in I_U$ be a Lipschitz function. As in the proof of Theorem 2.3, we have that $i_A(a) = \lim_{k \rightarrow \infty} \sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^*$. Let $\varepsilon > 0$. Since $\bigcap_{k \in \mathbf{N}} \phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)}) = \{x_\alpha\}$ with $x_\alpha \in K_{s(\alpha)}$ and $\text{diam}(\phi_{\alpha_1 \dots \alpha_k}(T_{r(\alpha_k)})) < c^k D$ for all $\alpha \in E^\infty$, there exists $N \in \mathbf{N}$ such that $|\mu_{r(\alpha_k)}^0(a \circ \phi_{\alpha_1 \dots \alpha_k}) - a(x_\alpha)| < \varepsilon$ for all $\alpha \in E^\infty$ and $k \geq N$. Since $a(x) = 0$ for all $x \in K$, $\|\sum_{\alpha \in E^k} \mu_{r(\alpha)}^0(a_{s(\alpha)} \circ \phi_\alpha) S_\alpha S_\alpha^*\| < \varepsilon$ for all $k \geq N$. Hence, $i_A(a) = 0$. \square

Corollary 2.6. *The Cuntz-Pimsner algebra associated to the C^* -correspondence $C(E \times_G K)$ over $C(K)$ with the actions defined as in Definition 2.2 is isomorphic to $C^*(G)$.*

One can interpret the previous results in the particular case of the iterated function system and obtain the result from [23, Remark 4.6].

3. On noncommutative Mauldin-Williams graphs. We give a generalization of the work of Pinzari, Watatani and Yonetani from [23, Section 4.3] on noncommutative iterated function systems in the context of “noncommutative” Mauldin-Williams graphs and the Rieffel metric. We show that, in fact, these situations are no more general than those just discussed.

We begin by reviewing the Rieffel metric.

Definition 3.1. Let A be a unital C^* -algebra, let $\mathcal{L}(A) \subset A$ be a dense subspace of A (the Lipschitz elements), and let L be a semi-norm (the Lipschitz semi-norm) on $\mathcal{L}(A)$ such that $\mathcal{K} := \{a \in \mathcal{L}(A) : L(a) = 0\}$ equals the scalar multiples of the identity. The Rieffel metric ρ on

the state space \mathcal{S} of A is defined by the equation

$$\rho(\mu, \nu) = \sup\{|\mu(a) - \nu(a)| : a \in \mathcal{L}(A), L(a) \leq 1\}$$

for all $\mu, \nu \in \mathcal{S}$. We will suppose that the metric ρ is bounded on \mathcal{S} and that the corresponding topology coincides with the weak-* topology on \mathcal{S} .

For a compact metric space (X, ρ) , let

$$\mathcal{C}(X) := \{E : E \text{ is a non-empty compact subset of } X\}.$$

The Hausdorff metric on $\mathcal{C}(X)$ is defined by the formula

$$\delta_\rho(E, F) = \inf\{r > 0 : U_r(E) \supseteq F \text{ and } U_r(F) \supseteq E\}$$

for all $E, F \in \mathcal{C}(X)$, where $U_r(E) = \{x \in X : \rho(x, y) < r \text{ for some } y \in E\}$, see [8, Theorem 2.4.1] or [15, Proposition 1.1.5]. Then $(\mathcal{C}(X), \delta_\rho)$ is a compact metric space.

For a C^* -algebra A , Rieffel defines the quantum closed subsets of A in [29, p. 14] to be the closed convex subsets of the state space $\mathcal{S}(A)$ of A . If L is a Lipschitz semi-norm on A and ρ_L is the corresponding Rieffel metric, the space $\mathcal{Q}(A)$ of quantum closed subsets of A is a compact metric space for the associated Hausdorff metric, see [29, p. 14]).

Following the definition of the classical Mauldin-Williams graphs, we define a noncommutative variant.

Definition 3.2. A noncommutative Mauldin-Williams graph is a system $(G, \{A_v, \mathcal{L}_v, L_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$ where $G = (V, E, s, r)$ is a graph and where $\{A_v, \mathcal{L}_v, L_v\}_{v \in V}$ and $\{\phi_e\}_{e \in E}$ are families such that

(1) For each $v \in V$, A_v is a unital C^* -algebra with a prescribed Lipschitz semi-norm L_v on a prescribed subspace \mathcal{L}_v of Lipschitz elements in A_v and ρ_v is the corresponding Rieffel metric.

(2) For $e \in E$, ϕ_e is a unital $*$ -homomorphism from $A_{s(e)}$ to $A_{r(e)}$ such that

$$\rho_{s(e)}(\phi_e^*(\mu), \phi_e^*(\nu)) \leq c\rho_{r(e)}(\mu, \nu)$$

for some constant c satisfying $0 < c < 1$ and all $\mu, \nu \in \mathcal{S}_{r(e)}$ (where \mathcal{S}_v is the state space of the C^* -algebra A_v).

We shall assume, too, that there are no sinks in the graph G . We also let $\mathcal{S} := \prod_{v \in V} \mathcal{S}_v$.

When we have one vertex and n edges we call the system a *noncommutative iterated function system*.

Let $\mathcal{C}(\mathcal{S}_v)$ be the space of compact subsets of \mathcal{S}_v endowed with the Hausdorff metric $\delta_{\rho_{L_v}}$, for each $v \in V$. Let $\mathcal{C} = \prod_{v \in V} \mathcal{C}(\mathcal{S}_v)$. Then \mathcal{C} is a compact metric space. Moreover, the map $F : \mathcal{C} \rightarrow \mathcal{C}$ defined by the formula

$$F((K_v)_{v \in V}) = \left(\bigcup_{\substack{e \in E \\ s(e)=v}} \phi_e^*(K_{r(e)}) \right)_{v \in V}$$

is a contraction, since each ϕ_e^* is a contraction with respect to the Rieffel metric. Thus, there exists a *unique* element $(K_v)_{v \in V} \in \mathcal{C}$ such that

$$(3.1) \quad K_v = \bigcup_{\substack{e \in E \\ s(e)=v}} \phi_e^*(K_{r(e)})$$

for all $v \in V$. Let T_v be the closed convex hull of K_v , for $v \in V$. That is $T_v \in \mathcal{Q}(A_v)$ for all $v \in V$. Since by [29, Proposition 3.6] there is a bijection between isomorphism classes of quotients of A_v and closed convex subsets of \mathcal{S}_v , we will assume that

$$(3.2) \quad \mathcal{S}_v = T_v \quad \text{for all } v \in V,$$

by taking a quotient of the original C^* -algebra A_v , if necessary. In particular, if $(M_v)_{v \in V} \in \mathcal{S}$ is any family which satisfies equation (3.1), then $M_v = K_v$ and the closed convex hull of M_v equals \mathcal{S}_v for all $v \in V$.

Lemma 3.3. *In the above situation, if \mathcal{I} is an ideal in $\sum_{v \in V}^\oplus A_v$ of the form $\mathcal{I} = (\mathcal{I}_v)_{v \in V}$, with \mathcal{I}_v a proper ideal of A_v , then*

$$(3.3) \quad \mathcal{I}_v = \bigcap_{\substack{e \in E \\ s(e)=v}} \phi_e^{-1}(\mathcal{I}_{r(e)}) \quad \text{if and only if} \quad \mathcal{I}_v = (0_v) \quad \text{for all } v \in V.$$

Proof. Let $\mathcal{I} = (\mathcal{I}_v)_{v \in V}$ be such that $\mathcal{I}_v = \bigcap_{e \in E, s(e)=v} \phi_e^{-1}(\mathcal{I}_{r(e)})$ for all $v \in V$. Let $M_v := \{\mu \in \mathcal{S}_v : \mu(a) = 0 \text{ for all } a \in \mathcal{I}_v\}$. We show that

$(M_v)_{v \in V}$ is a family which satisfies equation (3.1). Let $v \in V$, and let $\mu \in \cup_{e \in E, s(e)=v} \phi_e^*(M_{r(e)})$. Then there exists some $e \in E$ and $\nu \in M_{r(e)}$ such that $\mu = \phi_e^*(\nu)$. Let $a \in \mathcal{I}_v$. Then $\phi_e(a)$ belongs to $\mathcal{I}_{r(e)}$. Hence $\mu(a) = \nu(\phi_e(a)) = 0$. Therefore $\mu \in M_v$.

Now suppose that there is some $\mu \in M_v$ such that

$$\mu \notin \bigcup_{\substack{e \in E \\ s(e)=v}} \phi_e^*(M_{r(e)}).$$

Hence, there is some $a \in A_v$ such that $\mu(a) \neq 0$ and $\phi_e^*(\nu)(a) = 0$ for all $\nu \in M_{r(e)}$ and for all $e \in E$ such that $s(e) = v$. Then $\phi_e(a_v) \in \mathcal{I}_{r(e)}$ for all $e \in E$, therefore $a_v \in \mathcal{I}_v$. Thus $\mu(a) = 0$, which is a contradiction. Then the family $(M_v)_{v \in V}$ satisfies equation (3.1). Therefore $\mu(a) = 0$ for all $\mu \in \mathcal{S}_v$ and $a \in \mathcal{I}_v$, hence $\mathcal{I}_v = 0$ for all $v \in V$.

Suppose that there exists some $a \in A_v$ which is not zero, but $a \in \cap_{e \in E, s(e)=v} \text{Ker } \phi_e$. Then there is some $\mu \in \mathcal{S}_v$ such that $\mu(a) = 0$. Since $K_v = \cup_{e \in E, s(e)=v} \phi_e^*(K_{r(e)})$, there is some $e \in E$ with $s(e) = v$ and some $\nu \in \mathcal{S}_{r(e)}$ such that $\mu = \phi_e^*(\nu)$. Since $\phi_e(a) = 0$, we obtain that $\mu(a) = \nu(\phi_e(a)) = 0$, which is a contradiction. Hence $(0_v) = \cap_{e \in E, s(e)=v} \text{Ker } \phi_e$. \square

Recall that E^k denotes the set of paths of length k , E^∞ denotes the set of infinite paths in the graph G , $E^k(v)$ denotes the set of paths of length k starting at the vertex v , and $E^\infty(v)$ denotes the set of infinite paths starting at the vertex v . For $k \in \mathbf{N}$ and $\alpha \in E^k$, we write $\phi_{\alpha_1 \dots \alpha_k}^*$ for the map $\phi_{\alpha_1}^* \circ \dots \circ \phi_{\alpha_k}^* : \mathcal{S}_{r(\alpha_k)} \rightarrow \mathcal{S}_{s(\alpha_1)}$ and $\phi_{\alpha_k \dots \alpha_1}$ for the map $\phi_{\alpha_k} \circ \dots \circ \phi_{\alpha_1} : A_{s(\alpha_1)} \rightarrow A_{r(\alpha_k)}$. We will use the following results (which are similar to the commutative case): if $v \in V$, $a \in \mathcal{L}_v(A_v)$ and $\mu, \nu \in \mathcal{S}_v$ then $|\mu(a) - \nu(a)| \leq \rho_v(\mu, \nu) \cdot L_v(a)$; if $\alpha \in E^k$ and $\mu, \nu \in \mathcal{S}_{r(\alpha)}$, then

$$(3.4) \quad \rho_{s(\alpha)}(\phi_\alpha^*(\mu), \phi_\alpha^*(\nu)) \leq c^k \rho_{r(\alpha)}(\mu, \nu) \leq c^k D,$$

where $s(\alpha) = s(\alpha_1)$, $r(\alpha) = r(\alpha_k)$ and $D = \max_{v \in V} \text{diam}_{L_v}(\mathcal{S}_v)$.

Since $(G, \{\mathcal{S}_v, \rho_v\}_{v \in V}, \{\phi_e^*\}_{e \in E})$ is a (classical) Mauldin-Williams graph, for each $\alpha \in E^\infty$ there is a unique state $\mu_\alpha \in \mathcal{S}_{s(\alpha)}$ such that $\{\mu_\alpha\} = \cap_{k \in \mathbf{N}} \phi_{\alpha_1 \dots \alpha_k}^*(\mathcal{S}_{r(\alpha_k)})$. In particular $\lim_{k \rightarrow \infty} \phi_{\alpha_1 \dots \alpha_k}^*(\mu_{r(\alpha_k)}) = \mu_\alpha$ for all $\mu = (\mu_v)_{v \in V} \in \mathcal{S}$.

Theorem 3.4. *Let $(G, \{A_v, \mathcal{L}_v, L_v, \rho_v\}_{v \in V}, \{\phi_e\}_{e \in E})$ be a (non-commutative) Mauldin-Williams graph. Suppose that the graph G has no sinks. Then there is an injective $*$ -homomorphism from A into $C(E^\infty)$.*

Proof. Fix $v_0 \in V$. Define $\pi_{v_0} : A_{v_0} \rightarrow C(E^\infty(v_0))$ by the formula

$$\pi_{v_0}(a)(\alpha) = \mu_\alpha(a)$$

for all $a \in A_{v_0}$. Thus, if $a \in \mathcal{L}_{v_0}$,

$$\pi_{v_0}(a)(\alpha) = \lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}(\phi_{\alpha_k \cdots \alpha_1}(a))$$

for all $\mu = (\mu_v)_{v \in V} \in \mathcal{S}$. By the comments preceding the theorem the map π_{v_0} is well defined. We prove that it is a homomorphism.

Let $\mu_0 = (\mu_v^0)_{v \in V} \in \mathcal{S}$ be fixed. Let $a \in \mathcal{L}_{v_0}$, $\alpha \in E^\infty(v_0)$ and let $\varepsilon > 0$. Let $k \in \mathbf{N}$ be such that $c^k DL_{v_0}(a) < \varepsilon$. For any $\mu = (\mu_v)_{v \in V} \in \mathcal{S}$, we have

$$\begin{aligned} & \left| \mu_{r(\alpha_k)} \left(\mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(a)) 1_{A_{r(\alpha_k)}} - \phi_{\alpha_k \cdots \alpha_1}(a) \right) \right| \\ &= \left| \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(a)) - \mu_{r(\alpha_k)}(\phi_{\alpha_k \cdots \alpha_1}(a)) \right| < c^k DL_{v_0}(a) < \varepsilon. \end{aligned}$$

Hence,

$$(3.5) \quad \left\| \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(a)) 1_{A_{r(\alpha_k)}} - \phi_{\alpha_k \cdots \alpha_1}(a) \right\| < 4\varepsilon.$$

Let $a, b \in A_{v_0}$. We have

$$\begin{aligned} & \left| \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(ab)) - \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(a)) \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(b)) \right| \\ & \leq \left\| \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(ab)) - \phi_{\alpha_k \cdots \alpha_1}(a) \phi_{\alpha_k \cdots \alpha_1}(b) \right\| \\ & \quad + \left\| \phi_{\alpha_k \cdots \alpha_1}(a) \phi_{\alpha_k \cdots \alpha_1}(b) - \phi_{\alpha_k \cdots \alpha_1}(a) \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(b)) \right\| \\ & \quad + \left\| \phi_{\alpha_k \cdots \alpha_1}(a) \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(b)) \right. \\ & \quad \quad \left. - \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(a)) \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \cdots \alpha_1}(b)) \right\| \\ & < 4\varepsilon + 4\|a\|\varepsilon + 4\varepsilon = (8 + 4\|a\|)\varepsilon, \end{aligned}$$

by inequality (3.5). Since $\lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \dots \alpha_1}(ab)) = \pi_{v_0}(ab)(\alpha)$ and $\lim_{k \rightarrow \infty} \mu_{r(\alpha_k)}^0(\phi_{\alpha_k \dots \alpha_1}(a))\mu_{r(\alpha_k)}^0(\phi_{\alpha_k \dots \alpha_1}(b)) = \pi_{v_0}(a)(\alpha)\pi_{v_0}(b)(\alpha)$, we see that π_{v_0} is a homomorphism.

Hence, for each $v \in V$, we have defined an $*$ -homomorphism $\pi_v : A_v \rightarrow C(E^\infty(v))$. We prove that π_v is injective for all $v \in V$. Let $v \in V$. Let $a \in A_v$. Then

$$\begin{aligned} a \in \text{Ker } \pi_v &\iff \pi_v(a)(\alpha) = 0 \ \forall \ \alpha = (\alpha_n)_{n \in \mathbf{N}} \in E^\infty(v) \\ &\iff \pi_v(a)(e\beta) = 0 \ \forall \ \beta \in E^\infty(r(e)) \quad \text{and} \quad e \in E(v) \\ &\iff \pi_{r(e)}(\phi_e(a))(\beta) = 0 \ \forall \ \beta \in E^\infty(r(e)) \quad \text{and} \quad e \in E(v) \\ &\iff a \in \bigcap_{s(e)=v} \text{Ker } \pi_{r(e)} \circ \phi_e. \end{aligned}$$

Lemma 3.3 implies that $\text{Ker } \pi_v = 0$ for all $v \in V$, hence the $*$ -homomorphism $\pi : A \rightarrow C(E^\infty)$ defined by the formula

$$\pi((a_v)_{v \in V}) = \sum_{v \in V}^\oplus \pi_v(a_v)$$

is an injective $*$ -homomorphism. □

Corollary 3.5. *Under the hypothesis of Theorem 3.4, we conclude that A must be a commutative C^* -algebra.*

Even in the setting of a “noncommutative” iterated function system studied in [23, Section 4.2], if we have defined a Rieffel metric such that the underlying topology and the weak- $*$ topology coincide, and if the duals of the endomorphisms restricted to the state space of A are contractions with respect to the Rieffel metric, then (under the hypothesis that A satisfies equation (3.2)) A is forced to be commutative and the endomorphisms ϕ_i must come from an ordinary iterated function system, i.e., $A = C(K)$ for some compact metric space and there are contractions $\{\varphi_i\}_{i=1, \dots, n}$ defined on K such that $\phi_i(a) = a \circ \varphi_i$. This seems not to have been noticed by the authors of [23].

The assumption that the graph G has no sinks is essential in the proofs of Theorems 2.3 and 3.4, since it forces the presence of infinite paths in the graph. Also, the assumption that the graph G has no

sources was needed to define the C^* -correspondence associated with a Mauldin-Williams graph. It is not needed, though, in the proof of Theorem 3.4.

We would like to call attention to a recent preprint of Kajiwara and Watatani [13] in which they considered a somewhat different C^* -correspondences associated with an iterated function system and arrive at a C^* -algebra that is sometimes different from \mathcal{O}_n . It appears that their construction can be modified to cover the setting of Mauldin-Williams graphs, leading to C^* -algebras different from the Cuntz-Krieger algebras of the underlying graphs. We intend to pursue the ramifications of this in a future note.

Acknowledgments. The author would like to thank Mark Tomforde for many useful comments regarding the work on this paper and especially Paul Muhly for his constant encouragement and beneficial conversations. The author would also like to express his thanks to the referee for pointing out errors in the original draft of this note and for suggesting the use of the Gauge-Invariant Uniqueness Theorem in the proof of Theorem 2.3, and to Yasuo Watatani, Alex Kumjian, and Dorin Dutkay for useful conversations after the first version of the paper was written.

REFERENCES

1. M.F. Barnsley, *Fractals everywhere*, 2nd ed., Academic Press Professional, Boston, MA, 1993.
2. L. Bartholdi, R. Grigorchuk and V. Nekrashevych, *From fractal groups to fractal sets*, in *Trends in mathematics: Fractals in Graz 2001*, Birkhauser, Berlin, pp. 25–118.
3. J. Cuntz, *Simple C^* -algebras generated by isometries*, *Comm. Math. Phys.* **57** (1977), 173–185.
4. J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, *Invent. Math.* **56** (1980), 251–268.
5. V. Deaconu, *Groupoids associated with endomorphisms*, *Trans. Amer. Math. Soc.* **347** (1995), 1779–1786.
6. ———, *Generalized solenoids and C^* -algebras*, *Pacific J. Math.* **190** (1999), 247–260.
7. V. Deaconu, A. Kumjian and P. Muhly, *Cohomology of topological graphs and Cuntz-Pimsner algebras*, *J. Operator Theory* **46** (2001), 251–264.

8. G.A. Edgar, *Measure, topology, and fractal geometry*, Undergrad. Texts Math., Springer-Verlag, New York, 1990.
9. G.A. Edgar and R.D. Mauldin, *Multifractal decompositions of digraph recursive fractals*, Proc. London Math. Soc. **65** (1992), 604–628.
10. N.J. Fowler, M. Laca and I. Raeburn, *The C^* -algebras of infinite graphs*, Proc. Amer. Math. Soc. **128** (2000), 2319–2327.
11. N.J. Fowler, P.S. Muhly and I. Raeburn, *Representations of Cuntz-Pimsner algebras*, Indiana Univ. Math. J. **52** (2003), 569–605.
12. J.E. Hutchinson, *Fractals and self-similarity*, Indiana Univ. Math. J. **30** (1981), 713–747.
13. T. Kajiwara and Y. Watatani, *C^* -algebras associated with self-similar sets*, preprint OA/0312481.
14. T. Katsura, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras, I, Fundamental results*, Trans. Amer. Math. Soc. **356** (2004), 4287–4322.
15. J. Kigami, *Analysis on fractals*, Cambridge Univ. Press, Cambridge, 2001.
16. A. Kumjian, D. Pask and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), 161–174.
17. E.C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*, London Math. Soc. Lecture Note Ser., vol. 210, Cambridge Univ. Press, Cambridge, 1995.
18. R.D. Mauldin and S.C. Williams, *Hausdorff dimension in graph directed constructions*, Trans. Amer. Math. Soc. **309** (1988), 811–829.
19. P.S. Muhly and B. Solel, *On the Morita equivalence of tensor algebras*, Proc. London Math. Soc. **81** (2000), 113–168.
20. ———, *Tensor algebras over C^* -correspondences: Representations, dilations, and C^* -envelopes*, J. Funct. Anal. **158** (1998), 389–457.
21. P. Muhly and M. Tomforde, *Topological quivers*, Internat. J. Math. **16** (2005), 693–755.
22. M.V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbf{Z}* , in *Free probability theory* (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 189–212.
23. C. Pinzari, Y. Watatani and K. Yonetani, *KMS states, entropy and the variational principle in full C^* -dynamical systems*, Comm. Math. Phys. **213** (2000), 331–379.
24. I. Raeburn, *On the Picard group of a continuous trace C^* -algebra*, Trans. Amer. Math. Soc. **263** (1981), 183–205.
25. I. Raeburn and D. P. Williams, *Morita equivalence and continuous-trace C^* -algebras*, Math. Surveys Monogr., vol. 60, Amer. Math. Soc., Providence, 1998.
26. J. Renault, *Cuntz like-algebras*, J. Operator Theory **25** (1991), 3–36.
27. M.A. Rieffel, *Metrics on states from actions of compact groups*, Doc. Math. **3** (1998), 215–229.
28. ———, *Metrics on state spaces*, Doc. Math. **4** (1999), 559–600.

29. M.A. Rieffel, *Gromov-Hausdorff distance for quantum metric spaces*, Appendix 1 by Hanfeng Li. *Gromov-Hausdorff distance for quantum metric spaces. Matrix algebras converge to the sphere for quantum Gromov-Hausdorff distance*, Mem. Amer. Math. Soc. **168** (2004), 1–65.

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, 6188 KEMENY HALL,
HANOVER, NH 03755-3551

E-mail address: marius.ionescu@dartmouth.edu