

# SIMPLICITY OF $A_n$

KEITH CONRAD

## 1. INTRODUCTION

A finite group is called *simple* when its only normal subgroups are the trivial subgroup and the whole group.

For instance, a finite group of prime size is simple, since it in fact has no non-trivial proper subgroups at all (normal or not). A finite abelian group  $G$  not of prime size, is not simple: let  $p$  be a prime factor of  $\#G$ , so  $G$  contains a subgroup of order  $p$ , which is a normal since  $G$  is abelian and is proper since  $\#G > p$ . Thus, the abelian finite simple groups are the groups of prime size.

When  $n \geq 3$  the group  $S_n$  is not simple because it has a nontrivial normal subgroup  $A_n$ . But the groups  $A_n$  are simple, provided  $n \geq 5$ .

**Theorem 1.1** (C. Jordan, 1875). *For  $n \geq 5$ , the group  $A_n$  is simple.*

The restriction  $n \geq 5$  is optimal, since  $A_4$  is not simple: it has a normal subgroup of size 4, namely  $\{(1), (12)(34), (13)(24), (14)(23)\}$ . The group  $A_3$  is simple, since it has size 3, and the groups  $A_1$  and  $A_2$  are trivial.

We will give five proofs of Theorem 1.1. Section 2 includes some preparatory material and later sections give the proofs of Theorem 1.1. In the final section, we give a quick application of the simplicity of alternating groups and give references for further proofs not treated here.

## 2. PRELIMINARIES

We give two lemmas about alternating groups  $A_n$  for  $n \geq 5$  and then two results on symmetric groups  $S_n$  for  $n \geq 5$ .

**Lemma 2.1.** *For  $n \geq 3$ ,  $A_n$  is generated by 3-cycles. For  $n \geq 5$ ,  $A_n$  is generated by permutations of type  $(2, 2)$ .*

*Proof.* That the 3-cycles generate  $A_n$  for  $n \geq 3$  has been seen earlier in the course. To show permutations of type  $(2, 2)$  generate  $A_n$  for  $n \geq 5$ , it suffices to write any 3-cycle  $(abc)$  in terms of such permutations. Pick  $d, e \notin \{a, b, c\}$ . Then note

$$(abc) = (ab)(de)(de)(bc).$$

□

The 3-cycles in  $S_n$  are all conjugate in  $S_n$ , since permutations of the same cycle type in  $S_n$  are conjugate. Are 3-cycles conjugate in  $A_n$ ? Not when  $n = 4$ :  $(123)$  and  $(132)$  are not conjugate in  $A_4$ . But for  $n \geq 5$  we do have conjugacy in  $A_n$ .

**Lemma 2.2.** *For  $n \geq 5$ , any two 3-cycles in  $A_n$  are conjugate in  $A_n$ .*

*Proof.* We show every 3-cycle in  $A_n$  is conjugate within  $A_n$  to (123). Let  $\sigma$  be a 3-cycle in  $A_n$ . It can be conjugated to (123) in  $S_n$ :

$$(123) = \pi\sigma\pi^{-1}$$

for some  $\pi \in S_n$ . If  $\pi \in A_n$  we're done. Otherwise, let  $\pi' = (45)\pi$ , so  $\pi' \in A_n$  and

$$\pi'\sigma\pi'^{-1} = (45)\pi\sigma\pi^{-1}(45) = (45)(123)(45) = (123).$$

□

**Example 2.3.** The 3-cycles (123) and (132) are not conjugate in  $A_4$ . But in  $A_5$  we have

$$(132) = \pi(123)\pi^{-1}$$

for  $\pi = (45)(12) \in A_5$ .

Most proofs of the simplicity of the groups  $A_n$  are based on Lemmas 2.1 and 2.2. The basic argument is this: show any non-trivial normal subgroup  $N \triangleleft A_n$  contains a 3-cycle, so  $N$  contains every 3-cycle by Lemma 2.2, and therefore  $N$  is  $A_n$  by Lemma 2.1.

The next lemma will be used in our fifth proof of the simplicity of alternating groups.

**Lemma 2.4.** *For  $n \geq 5$ , the only nontrivial proper normal subgroup of  $S_n$  is  $A_n$ . In particular, the only subgroup of  $S_n$  with index 2 is  $A_n$ .*

*Proof.* The last statement follows from the first since any subgroup of index 2 is normal.

Let  $N \triangleleft S_n$  with  $N \neq \{(1)\}$ . We will show  $A_n \subset N$ , so  $N = A_n$  or  $S_n$ .

Pick  $\sigma \in N$  with  $\sigma \neq (1)$ . That means there is an  $i$  with  $\sigma(i) \neq i$ . Pick  $j \in \{1, 2, \dots, n\}$  so  $j \neq i$  and  $j \neq \sigma(i)$ . Let  $\tau = (ij)$ . Then

$$\sigma\tau\sigma^{-1}\tau^{-1} = (\sigma(i) \sigma(j))(ij).$$

Since  $\sigma(i) \neq i$  or  $j$  and  $\sigma(i) \neq \sigma(j)$  (why?), the 2-cycles  $(\sigma(i) \sigma(j))$  and  $(ij)$  are unequal, so their product is not the identity. That shows  $\sigma\tau \neq \tau\sigma$ .

Since  $N \triangleleft S_n$ ,  $\sigma\tau\sigma^{-1}\tau^{-1}$  lies in  $N$ . By construction,  $\sigma(i) \neq i$  or  $j$ . If  $\sigma(j) \neq i$  or  $j$ , then  $(\sigma(i) \sigma(j))(ij)$  has type (2, 2). If  $\sigma(j) = i$  or  $j$ ,  $(\sigma(i) \sigma(j))(ij)$  is a 3-cycle. Thus  $N$  contains a permutation of type (2, 2) or a 3-cycle. Since  $N \triangleleft S_n$ ,  $N$  contains all permutations of type (2, 2) or all 3-cycles. In either case, this shows (by Lemma 2.1) that  $N \supset A_n$ . □

**Remark 2.5.** There is an analogue of Lemma 2.4 for the “countable” symmetric group  $S_\infty$  consisting of all permutations of  $\{1, 2, 3, \dots\}$ . A theorem of Schreier and Ulam (1933) says the only nontrivial proper normal subgroups of  $S_\infty$  are  $\cup_{n \geq 1} S_n$  and  $\cup_{n \geq 1} A_n$ , which are the subgroup of permutations fixing all but a finite number of terms and its subgroup of even permutations.

**Remark 2.6.** From Lemma 2.4, any homomorphic image of  $S_n$  which is not an isomorphism has size 1 or 2. In particular, there is no surjective homomorphism  $S_n \rightarrow \mathbf{Z}/(m)$  for  $m > 2$ .

**Theorem 2.7.** *For  $n \geq 5$ , any proper subgroup of  $S_n$  other than  $A_n$  has index at least  $n$ . Moreover, any subgroup of index  $n$  is isomorphic to  $S_{n-1}$ .*

*Proof.* Let  $H$  be a proper subgroup of  $S_n$  other than  $A_n$ , and let  $m > 1$  be the index of  $H$  in  $S_n$ . We want to show  $m \geq n$ . Assume  $m < n$ . The left multiplication action of  $S_n$  on  $S_n/H$  gives a group homomorphism

$$\varphi: S_n \rightarrow \text{Sym}(S_n/H) \cong S_m.$$

By hypothesis,  $m < n$ , so  $\varphi$  is not injective. Let  $K$  be the kernel of  $\varphi$ , so  $K \subset H$  and  $K$  is non-trivial. Since  $K \triangleleft S_n$ , Lemma 2.4 says  $K = A_n$  or  $S_n$ . Since  $K \subset H$ , we get  $H = A_n$  or  $S_n$ , which contradicts our initial assumption about  $H$ . Therefore  $m \geq n$ .

Now let  $H$  be a subgroup of  $S_n$  with index  $n$ . Consider the left multiplication action of  $S_n$  on  $S_n/H$ . This is a homomorphism  $\ell: S_n \rightarrow \text{Sym}(S_n/H)$ . Since  $S_n/H$  has size  $n$ ,  $\text{Sym}(S_n/H)$  is isomorphic to  $S_n$ . The kernel of  $\ell$  is a normal subgroup of  $S_n$  which lies in  $H$  (why?). Therefore the kernel has index at least  $n$  in  $S_n$ . Since the only normal subgroups of  $S_n$  are 1,  $A_n$ , and  $S_n$ , the kernel of  $\ell$  is trivial, so  $\ell$  is an isomorphism. What is the image  $\ell(H)$  in  $\text{Sym}(S_n/H)$ ? Since  $gH = H$  if and only if  $g \in H$ ,  $\ell(H)$  is the group of permutations of  $S_n/H$  which fixes the ‘‘point’’  $H$  in  $S_n/H$ . The subgroup fixing a point in a symmetric group isomorphic to  $S_n$  is isomorphic to  $S_{n-1}$ . Therefore  $H \cong \ell(H) \cong S_{n-1}$ .  $\square$

Theorem 2.7 is false for  $n = 4$ :  $S_4$  contains the dihedral group of size 8 as a subgroup of index 3. An analogue of Theorem 2.7 for alternating groups will be given in Section 8; its proof uses the simplicity of alternating groups.

**Corollary 2.8.** *Let  $F$  be a field. If  $f \in F[X_1, \dots, X_n]$  and  $n \geq 5$ , the number of different polynomials we get from  $f$  by permuting its variables is either 1, 2, or at least  $n$ .*

*Proof.* Letting  $S_n$  act on  $F[X_1, \dots, X_n]$  by permutations of the variables, the polynomials we get by permuting the variables of  $f$  is the  $S_n$ -orbit of  $f$ . The size of this orbit is  $[S_n : H]$ , where  $H = \text{Stab}_f = \{\sigma \in S_n : \sigma f = f\}$ . By Theorem 2.7, this index is either 1, 2, or at least  $n$ .  $\square$

### 3. FIRST PROOF

Our first proof of Theorem 1.1 is based on the one in [2, pp. 149–150].

We begin by showing  $A_5$  is simple.

**Theorem 3.1.** *The group  $A_5$  is simple.*

*Proof.* We want to show the only normal subgroups of  $A_5$  are  $\{(1)\}$  and  $A_5$ . This will be done in two ways.

Our first method involves counting the sizes of the conjugacy classes. There are 5 conjugacy classes in  $A_5$ , with representatives and sizes as indicated in the following table.

Rep.	(1)	(12345)	(21345)	(12)(34)	(123)
Size	1	12	12	15	20

If  $A_5$  has a normal subgroup  $N$ , then  $N$  is a union of conjugacy classes – including  $\{(1)\}$  – whose total size divides 60. However, no sum of the above numbers which includes 1 is a factor of 60 except for 1 and 60. Therefore  $N$  is trivial or  $A_5$ .

For the second proof, let  $N \triangleleft A_5$  with  $\#N > 1$ . We will show  $N$  contains a 3-cycle. It follows that  $N = A_n$  by Lemmas 2.1 and 2.2.

Pick  $\sigma \in N$  with  $\sigma \neq (1)$ . The cycle structure of  $\sigma$  is  $(abc)$ ,  $(ab)(cd)$ , or  $(abcde)$ , where different letters represent different numbers. Since we want to show  $N$  contains a 3-cycle, we may suppose  $\sigma$  has the second or third cycle type. In the second case,  $N$  contains

$$((abe)(ab)(cd)(abe)^{-1})(ab)(cd) = (be)(cd)(ab)(cd) = (aeb).$$

In the third case,  $N$  contains

$$((abc)(abcde)(abc)^{-1})(abcde)^{-1} = (adebc)(aedcb) = (abd).$$

Therefore  $N$  contains a 3-cycle, so  $N = A_5$ .  $\square$

**Lemma 3.2.** *When  $n \geq 5$ , any  $\sigma \neq (1)$  in  $A_n$  has a conjugate  $\sigma' \neq \sigma$  such that  $\sigma(i) = \sigma'(i)$  for some  $i$ .*

For example, if  $\sigma = (12345)$  in  $A_5$  then  $\sigma' = (345)\sigma(345)^{-1} = (12453)$  has the same value at  $i = 1$  as  $\sigma$  does.

*Proof.* Let  $\sigma$  be a non-identity element of  $A_n$ . Let  $r$  be the longest length of a disjoint cycle in  $\sigma$ . Relabelling, we may write

$$\sigma = (12 \dots r)\pi,$$

where  $(12 \dots r)$  and  $\pi$  are disjoint.

If  $r \geq 3$ , let  $\tau = (345)$  and  $\sigma' = \tau\sigma\tau^{-1}$ . Then  $\sigma(1) = 2, \sigma'(1) = 2, \sigma(2) = 3$ , and  $\sigma'(2) = 4$ . Thus  $\sigma' \neq \sigma$  and both take the same value at 1.

If  $r = 2$ , then  $\sigma$  is a product of disjoint transpositions. If there are at least 3 disjoint transpositions involved, then  $n \geq 6$  and we can write  $\sigma = (12)(34)(56)(\dots)$  after relabelling. Let  $\tau = (12)(35)$  and  $\sigma' = \tau\sigma\tau^{-1}$ . Then  $\sigma(1) = 2, \sigma'(1) = 2, \sigma(3) = 4$ , and  $\sigma'(3) = 6$ . Again, we see  $\sigma' \neq \sigma$  and  $\sigma$  and  $\sigma'$  have the same value at 1.

If  $r = 2$  and  $\sigma$  is a product of 2 disjoint transpositions, write  $\sigma = (12)(34)$  after relabelling. Let  $\tau = (132)$  and  $\sigma' = \tau\sigma\tau^{-1} = (13)(24)$ . Then  $\sigma' \neq \sigma$  and they both fix 5.  $\square$

Now we prove Theorem 1.1.

*Proof.* We may suppose  $n \geq 6$ , by Theorem 3.1. For  $1 \leq i \leq n$ , let  $A_n$  act in the natural way on  $\{1, 2, \dots, n\}$  and let  $H_i \subset A_n$  be the subgroup fixing  $i$ , so  $H_i \cong A_{n-1}$ . By induction, each  $H_i$  is simple. Note each  $H_i$  contains a 3-cycle (build out of 3 numbers other than  $i$ ).

Let  $N \triangleleft A_n$  be a nontrivial normal subgroup. We want to show  $N = A_n$ . Pick  $\sigma \in N$  with  $\sigma \neq \{(1)\}$ . By Lemma 3.2, there is a conjugate  $\sigma'$  of  $\sigma$  such that  $\sigma' \neq \sigma$  and  $\sigma(i) = \sigma'(i)$  for some  $i$ . Since  $N$  is normal in  $A_n$ ,  $\sigma' \in N$ . Then  $\sigma^{-1}\sigma'$  is a non-identity element of  $N$  which fixes  $i$ , so  $N \cap H_i$  is a non-trivial subgroup of  $H_i$ . It is also a normal subgroup of  $H_i$  since  $N \triangleleft A_n$ . Since  $H_i$  is simple,  $N \cap H_i = H_i$ . Therefore  $H_i \subset N$ . Since  $H_i$  contains a 3-cycle,  $N$  contains a 3-cycle and we are done.

Alternatively, we can show  $N = A_n$  when  $N \cap H_i$  is non-trivial for some  $i$  as follows. As before, since  $N \cap H_i$  is a non-trivial normal subgroup of  $H_i$ ,  $H_i \subset N$ . Without referring to 3-cycles, we instead note that the different  $H_i$ 's are conjugate subgroups of  $A_n$ :  $\sigma H_i \sigma^{-1} = H_{\sigma(i)}$  for  $\sigma \in A_n$ . Since  $N \triangleleft A_n$  and  $N$  contains  $H_i$ ,  $N$  contains every  $H_{\sigma(i)}$  for all  $\sigma \in A_n$ . Since  $\sigma(i)$  can be any element of  $A_n$  as  $\sigma$  varies in  $A_n$ ,  $N$  contains every  $H_i$ . Any permutation of type  $(2, 2)$  is in some  $H_i$  since  $n \geq 5$ , so  $N$  contains all permutations of type  $(2, 2)$ . Every permutation in  $A_n$  is a product of permutations of type  $(2, 2)$ , so  $N \supset A_n$ . Therefore  $N = A_n$ .  $\square$

#### 4. SECOND PROOF

Our next proof is taken from [6, p. 108]. It does not use induction on  $n$ , but we do need to know  $A_6$  is simple at the start.

**Theorem 4.1.** *The group  $A_6$  is simple.*

*Proof.* We follow the first method of proof of Theorem 3.1. Here is the table of conjugacy classes in  $A_6$ .

Rep.	(1)	(123)	(123)(456)	(12)(34)	(12345)	(23456)	(1234)(56)
Size	1	40	40	45	72	72	90

A tedious check shows no sum of these sizes, which includes 1, is a factor of  $6!/2$  except for the sum of all the terms. Therefore the only non-trivial normal subgroup of  $A_6$  is  $A_6$ .  $\square$

Now we prove the simplicity of  $A_n$  for larger  $n$  by reducing directly to the case of  $A_6$ .

*Proof.* Since  $A_5$  and  $A_6$  are known to be simple by Theorems 3.1 and 4.1, pick  $n \geq 7$  and let  $N \triangleleft A_n$  be a non-trivial subgroup. We will show  $N$  contains a 3-cycle.

Let  $\sigma$  be a non-identity element of  $N$ . It moves some number. By relabelling, we may suppose  $\sigma(1) \neq 1$ . Let  $\tau = (ijk)$ , where  $i, j, k$  are not 1 and  $\sigma(1) \in \{i, j, k\}$ . Then  $\tau\sigma\tau^{-1}(1) = \tau(\sigma(1)) \neq \sigma(1)$ , so  $\tau\sigma\tau^{-1} \neq \sigma$ . Let  $\varphi = \tau\sigma\tau^{-1}\sigma^{-1}$ , so  $\varphi \neq (1)$ . Writing

$$\varphi = (\tau\sigma\tau^{-1})\sigma^{-1},$$

we see  $\varphi \in N$ . Now write

$$\varphi = \tau(\sigma\tau^{-1}\sigma^{-1}),$$

Since  $\tau^{-1}$  is a 3-cycle,  $\sigma\tau^{-1}\sigma^{-1}$  is also a 3-cycle. Therefore  $\varphi$  is a product of two 3-cycles, so  $\varphi$  moves at most 6 numbers in  $\{1, 2, \dots, n\}$ . Let  $H$  be the copy of  $A_6$  inside  $A_n$  corresponding to the even permutations of those 6 numbers (possibly augmented to 6 arbitrarily if in fact  $\varphi$  moves fewer numbers). Then  $N \cap H$  is non-trivial (it contains  $\varphi$ ) and it is a normal subgroup of  $H$ . Since  $H \cong A_6$ , which is simple,  $N \cap H = H$ . Thus  $H \subset N$ , so  $N$  contains a 3-cycle.  $\square$

## 5. THIRD PROOF

Our next proof is by induction, and uses conjugacy classes instead of Lemma 3.2. It is based on [9, §2.3].

**Lemma 5.1.** *If  $n \geq 6$  then every non-trivial conjugacy class in  $S_n$  and  $A_n$  has at least  $n$  elements.*

The lower bound  $n$  in Lemma 5.1 is actually quite weak as  $n$  grows. But it shows that the size of each non-trivial conjugacy class in  $S_n$  and  $A_n$  grows with  $n$ .

*Proof.* For  $n \geq 6$ , pick  $\sigma \in S_n$  with  $\sigma \neq (1)$ . We want to look at the conjugacy class of  $\sigma$  in  $S_n$ , and if  $\sigma \in A_n$  we also want to look at the conjugacy class of  $\sigma$  in  $A_n$ , and our goal in both cases is to find at least  $n$  elements in the conjugacy class.

**Case 1:** The disjoint cycle decomposition of  $\sigma$  includes a cycle with length greater than 2. Without loss of generality,  $\sigma = (123\dots)\dots$ .

For  $3 \leq k \leq n$ , fix a choice of  $\ell \notin \{1, 2, 3, k\}$  (which is possible since  $n \geq 5$ ) and let  $\alpha_k = (2k\ell)$  and  $\beta_k = (3k\ell)$ . Then  $\alpha_k\sigma\alpha_k^{-1}$  has the effect  $1 \rightarrow 1 \rightarrow 2 \rightarrow k$  and  $\beta_k\sigma\beta_k^{-1}$  has the effect  $1 \rightarrow 1 \rightarrow 2 \rightarrow 2$  and  $2 \rightarrow 2 \rightarrow 3 \rightarrow k$ . This tells us that the conjugates

$$\alpha_3\sigma\alpha_3^{-1}, \dots, \alpha_n\sigma\alpha_n^{-1}, \beta_3\sigma\beta_3^{-1}, \dots, \beta_n\sigma\beta_n^{-1}$$

are all different from each other: the conjugates by the  $\alpha$ 's have different effects on 1, the conjugates by the  $\beta$ 's have different effects on 2, and a conjugate by an  $\alpha$  is not a conjugate by a  $\beta$  since they have different effects on 1. Since these conjugates are different, the number of conjugates of  $\sigma$  is at least  $2(n-2) > n$ . Because  $\alpha_k$  and  $\beta_k$  are 3-cycles, if  $\sigma \in A_n$  then these conjugates are in the  $A_n$ -conjugacy class of  $\sigma$ .

**Case 2:** The disjoint cycle decomposition of  $\sigma$  only has cycles with length 1 or 2. Therefore without loss of generality  $\sigma$  is a transposition or a product of at least 2 disjoint transpositions.

If  $\sigma$  is a transposition, then its  $S_n$ -conjugacy class is the set of all transpositions  $(ij)$  where  $1 \leq i < j \leq n$ , and the number of these permutations is  $\binom{n}{2} = \frac{n^2-n}{2}$ , which is greater than  $n$  for  $n \geq 6$ .

If  $\sigma$  is a product of at least 2 disjoint transpositions, then without loss of generality  $\sigma = (12)(34)\dots$ , where the terms in  $\dots$  don't involve 1, 2, 3, or 4.

For  $5 \leq k \leq n$ , let  $\alpha_k = (12)(3k)$ ,  $\beta_k = (13)(2k)$ , and  $\gamma_k = (1k)(23)$ . Then  $\alpha_k \sigma \alpha_k^{-1}$  has the effect

$$1 \rightarrow 2 \rightarrow 1 \rightarrow 2, \quad 2 \rightarrow 1 \rightarrow 2 \rightarrow 1, \quad k \rightarrow 3 \rightarrow 4 \rightarrow 4,$$

$\beta_k \sigma \beta_k^{-1}$  has the effect

$$1 \rightarrow 3 \rightarrow 4 \rightarrow 4, \quad 3 \rightarrow 1 \rightarrow 2 \rightarrow k, \quad k \rightarrow 2 \rightarrow 1 \rightarrow 3,$$

and  $\gamma_k \sigma \gamma_k^{-1}$  has the effect

$$2 \rightarrow 3 \rightarrow 4 \rightarrow 4, \quad 3 \rightarrow 2 \rightarrow 1 \rightarrow k, \quad k \rightarrow 1 \rightarrow 2 \rightarrow 3.$$

The conjugates of  $\sigma$  by the  $\alpha$ 's are different from each other since they take different elements to 4, the conjugates of  $\sigma$  by the  $\beta$ 's are different from each other since they take different elements to 3, and the conjugates of  $\sigma$  by the  $\gamma$ 's are different from each other since they take different elements to 3. Conjugates of  $\sigma$  by an  $\alpha$  and a  $\beta$  are different since they send 1 to different places, conjugates of  $\sigma$  by an  $\alpha$  and a  $\gamma$  are different since they send 2 to different places, and conjugates of  $\sigma$  by a  $\beta$  and a  $\gamma$  are different since they send different elements to 4 (1 for the  $\beta$ 's and 2 for the  $\gamma$ 's). In total the number of conjugates of  $\sigma$  we have written down (which are all conjugates by 3-cycles, hence they are conjugates in  $A_n$  if  $\sigma \in A_n$ ) is  $3(n-4)$ , and  $3(n-4) \geq n$  if  $n \geq 6$ .  $\square$

Now we prove Theorem 1.1.

*Proof.* We argue by induction on  $n$ , the case  $n = 5$  having already been settled by Theorem 3.1. Say  $n \geq 6$ . Let  $N \triangleleft A_n$  with  $N \neq \{(1)\}$ . Since  $N$  is normal and non-trivial, it contains non-identity conjugacy classes in  $A_n$ . By Lemma 5.1, any non-identity conjugacy class in  $A_n$  has size at least  $n$  when  $n \geq 6$ . Therefore, by counting the trivial conjugacy class and a non-trivial conjugacy class in  $N$ , we see  $\#N \geq n + 1$ .

Using a wholly different argument, we now show that  $\#N \leq n$  if  $N \neq A_n$ , which will be a contradiction. Pick  $1 \leq i \leq n$ . Let  $H_i \subset A_n$  be the subgroup fixing  $i$ , so  $H_i \cong A_{n-1}$ . In particular,  $H_i$  is a simple group by induction. Notice each  $H_i$  contains a 3-cycle.

The intersection  $N \cap H_i$  is a normal subgroup of  $H_i$ , so simplicity of  $H_i$  implies  $N \cap H_i$  is either  $\{(1)\}$  or  $H_i$ . If  $N \cap H_i = H_i$  for some  $i$ , then  $H_i \subset N$ . Since  $H_i$  contains a 3-cycle,  $N$  does as well, so  $N = A_n$  by Lemmas 2.1 and 2.2. (This part resembles part of our first proof of simplicity of  $A_n$ , but we will now use Lemma 5.1 instead of Lemma 3.2 to show the possibility that  $N \cap H_i = \{(1)\}$  for all  $i$  is absurd.)

Suppose  $N \neq A_n$ . Then, by the previous paragraph,  $N \cap H_i = \{(1)\}$  for all  $i$ . Therefore each  $\sigma \neq (1)$  in  $N$  acts on  $\{1, 2, \dots, n\}$  without fixed points (otherwise  $\sigma$  would be a non-identity element in some  $N \cap H_i$ ). That implies each  $\sigma \neq (1)$  in  $N$  is completely determined by the value  $\sigma(1)$ : if  $\tau \neq (1)$  is in  $N$  and  $\sigma(1) = \tau(1)$ , then  $\sigma\tau^{-1} \in N$  fixes 1, so  $\sigma\tau^{-1}$  is the identity, so  $\sigma = \tau$ .

There are only  $n - 1$  possible values for  $\sigma(1) \in \{2, 3, \dots, n\}$ , so  $N - \{(1)\}$  has size at most  $n - 1$ , hence  $\#N \leq n$ . We already saw from Lemma 5.1 that  $\#N \geq n + 1$ , so we have a contradiction.  $\square$

## 6. FOURTH PROOF

Our next proof, based on [3, p. 50], is very computational.

*Proof.* Let  $N \triangleleft A_n$  be a non-trivial normal subgroup. We will show  $N$  contains a 3-cycle.

Pick  $\sigma \in N$ ,  $\sigma \neq (1)$ . Write

$$\sigma = \pi_1 \pi_2 \cdots \pi_k,$$

where the  $\pi_j$ 's are disjoint cycles. In particular, they *commute*, so we can relabel them at our convenience. Eliminate any 1-cycles from the product.

Case 1: Some  $\pi_i$  has length at least 4. Relabelling, we can write

$$\pi_1 = (12 \cdots r)$$

with  $r \geq 4$ . Let  $\varphi = (123)$ . Then  $\varphi \sigma \varphi^{-1} \in N$  and

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \varphi^{-1} \pi_2 \cdots \pi_k \\ &= \varphi \pi_1 \varphi^{-1} \pi_1^{-1} \sigma \\ &= (123)(123 \cdots r)(132)(r \cdots 21)\sigma \\ &= (124)\sigma, \end{aligned}$$

so  $(124) = \varphi \sigma \varphi^{-1} \sigma^{-1} \in N$ .

Case 2: Each  $\pi_i$  has length  $\leq 3$ , and at least two have length 3 (so  $n \geq 6$ ). Without loss of generality,  $\pi_1 = (123)$  and  $\pi_2 = (456)$ . Let  $\varphi = (124)$ . Then

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_3 \cdots \pi_k \\ &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_2^{-1} \pi_1^{-1} \sigma \\ &= (124)(123)(456)(142)(465)(132)\sigma \\ &= (12534)\sigma, \end{aligned}$$

so  $\varphi \sigma \varphi^{-1} \sigma^{-1} = (12534) \in N$ . Now run through Case 1 with this 5-cycle to find a 3-cycle in  $N$ .

Case 3: Exactly one  $\pi_i$  has length 3, and the rest have length  $\leq 2$ . Without loss of generality,  $\pi_1 = (123)$  and the other  $\pi_i$ 's are 2-cycles. Then  $\sigma^2 = \pi_1^2$  is in  $N$ , and  $\pi_1^2 = (132)$ .

Case 4: All  $\pi_i$ 's are 2-cycles, so necessarily  $k > 1$ . Write  $\pi_1 = (12)$  and  $\pi_2 = (34)$ . Let  $\varphi = (123)$ . Then

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_3 \cdots \pi_k \\ &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_2^{-1} \pi_1^{-1} \sigma \\ &= (123)(12)(34)(132)(34)(12)\sigma \\ &= (13)(24)\sigma, \end{aligned}$$

so

$$\varphi \sigma \varphi^{-1} \sigma^{-1} = (13)(24) \in N.$$

Let  $\psi = (135)$ . Then

$$\begin{aligned} (13)(24)\psi(13)(24)\psi^{-1} &= (13)(24)(135)(13)(24)(153) \\ &= (13)(135)(13)(153) \\ &= (135), \end{aligned}$$

so  $N$  contains a 3-cycle. □

## 7. FIFTH PROOF

Our final proof is taken from [8, p. 295].

Let  $N \triangleleft A_n$  with  $N$  not  $\{(1)\}$  or  $A_n$ . We will study  $N$  as a subgroup of  $S_n$ . By Lemma 2.4,  $N$  is not a normal subgroup of  $S_n$ . This means the normalizer of  $N$  inside  $S_n$  is a proper subgroup, which contains  $A_n$ , so

$$(7.1) \quad A_n = N_{S_n}(N).$$

For any transposition  $\tau$  in  $S_n$ ,  $\tau \notin N_{S_n}(N)$  by (7.1), so  $\tau N \tau^{-1} \neq N$ . Since  $N \triangleleft A_n$  and  $\tau N \tau^{-1}$  is a subgroup of  $A_n$ , the product set  $N \cdot \tau N \tau^{-1}$  is a subgroup of  $A_n$ . We have the chain of inclusions

$$N \cap \tau N \tau^{-1} \subset N \subset N \cdot \tau N \tau^{-1} \subset A_n,$$

where the first and second are strict.

We will now show, for any transposition  $\tau$  in  $S_n$ , that

$$(7.2) \quad N \cap \tau N \tau^{-1} \triangleleft S_n, \quad N \cdot \tau N \tau^{-1} \triangleleft S_n.$$

The proof of (7.2) is a bit tedious, so first let's see why (7.2) leads to a contradiction.

It follows from (7.2) and Lemma 2.4 that

$$(7.3) \quad N \cap \tau N \tau^{-1} = \{(1)\}, \quad N \cdot \tau N \tau^{-1} = A_n$$

for any transposition  $\tau$  in  $S_n$ . By (7.3),  $\#A_n = \#N \cdot \#(\tau N \tau^{-1}) = (\#N)^2$ , so  $n! = 2(\#N)^2$ . This tells us  $\#N$  must be even, so  $N$  has an element, say  $\sigma$ , of order 2. Then  $\sigma$  is a product of disjoint 2-cycles. There is a transposition  $\rho$  in  $S_n$  which commutes with  $\sigma$ : just take for  $\rho$  one of the transpositions in the disjoint cycle decomposition of  $\sigma$ . Then

$$\sigma = \rho \sigma \rho^{-1} \in N \cap \rho N \rho^{-1}.$$

From (7.3), using  $\rho$  for the arbitrary  $\tau$  there,  $N \cap \rho N \rho^{-1}$  is trivial, so we have a contradiction. (As another way of reaching a contradiction from the equation  $n! = 2(\#N)^2$ , we can use Bertrand's postulate – proved by Chebyshev – that there is always a prime strictly between  $m$  and  $2m$  for any  $m > 1$ . That means, taking  $m = n!/4$ , the ratio  $n!/2$  can't be a perfect square.)

It remains to check the two conditions in (7.2). In both cases, we show the subgroups are normalized by  $A_n$  and by  $\tau$ , so the normalizer contains  $\langle A_n, \tau \rangle = S_n$ .

First consider  $N \cap \tau N \tau^{-1}$ . It is clearly normalized by  $\tau$ . Now pick any  $\pi \in A_n$ . Then  $\pi N \pi^{-1} = N$  since  $N \triangleleft A_n$ , and

$$(7.4) \quad \pi(\tau N \tau^{-1})\pi^{-1} = \tau(\tau^{-1}\pi\tau)N(\tau^{-1}\pi^{-1}\tau)\tau^{-1} = \tau N \tau^{-1}$$

since  $\tau^{-1}\pi\tau \in A_n$ . Therefore

$$\pi(N \cap \tau N \tau^{-1})\pi^{-1} = \pi N \pi^{-1} \cap \pi \tau N \tau^{-1} \pi^{-1} = N \cap \tau N \tau^{-1},$$

so  $A_n$  normalizes  $N \cap \tau N \tau^{-1}$ .

Now we look at  $N \cdot \tau N \tau^{-1}$ . Pick an element of this product, say

$$\sigma = \sigma_1 \tau \sigma_2 \tau^{-1},$$

where  $\sigma_1, \sigma_2 \in N$ . Then, since  $N \triangleleft A_n$ ,

$$\tau \sigma \tau^{-1} = \tau \sigma_1 \tau \sigma_2 \tau^{-2} = \tau \sigma_1 \tau \sigma_2 \in \tau N \tau^{-1} \cdot N = N \cdot \tau N \tau^{-1},$$

which shows  $\tau$  normalizes  $N \cdot \tau N \tau^{-1}$ .



Now pick any  $\pi \in A_n$ . To see  $\pi$  normalizes  $N \cdot \tau N \tau^{-1}$ , pick  $\sigma$  as before. Then

$$\pi \sigma \pi^{-1} = \pi \sigma_1 \pi^{-1} \cdot \pi (\tau \sigma_2 \tau^{-1}) \pi^{-1}.$$

The first factor  $\pi \sigma_1 \pi^{-1}$  is in  $N$  since  $N \triangleleft A_n$ . The second factor is in  $\pi \tau N \tau^{-1} \pi^{-1}$ , which equals  $\tau N \tau^{-1}$  by (7.4).

## 8. CONCLUDING REMARKS

The standard counterexample to the converse of Lagrange's theorem is  $A_4$ : it has size 12 but no subgroup of index 2. For  $n \geq 5$ , the groups  $A_n$  also have no subgroup of index 2, since any index-2 subgroup of a group is normal and  $A_n$  is simple.

In fact, something stronger is true.

**Corollary 8.1.** *For  $n \geq 5$ , any proper subgroup of  $A_n$  has index at least  $n$ .*

This is an analogue of Theorem 2.7.

*Proof.* Let  $H$  be a proper subgroup of  $A_n$ , with index  $m > 1$ . Consider the left multiplication action of  $A_n$  on  $A_n/H$ . This gives a group homomorphism

$$\varphi: A_n \rightarrow \text{Sym}(A_n/H) \cong S_m.$$

Let  $K$  be the kernel of  $\varphi$ , so  $K \subset H$  (why?) and  $K \triangleleft A_n$ . By simplicity of  $A_n$ ,  $K$  is trivial. Therefore  $A_n$  injects into  $S_m$ , so  $(n!/2) | m!$ , which implies  $n \leq m$ .  $\square$

The lower bound of  $n$  is sharp since  $[A_n : A_{n-1}] = n$ . Corollary 8.1 is false for  $n = 4$ :  $A_4$  has a subgroup of index 3.

**Remark 8.2.** What the proof of Corollary 8.1 shows more generally is that if  $G$  is a finite simple group and  $H$  is a subgroup with index  $m > 1$ , then there is an embedding of  $G$  into  $S_m$ , so  $\#G | m!$ . With  $G$  fixed, this divisibility relation puts a lower bound on the index of any proper subgroup of  $G$ .

A reader who wants to see more proofs that  $A_n$  is simple for  $n \geq 5$  can look at [4, pp. 247–248] or [5, pp. 32–33] for another way of showing a non-trivial normal subgroup contains a 3-cycle, or at [1, §1.7] or [7, pp. 295–296] for a proof based on the theory of highly transitive permutation groups.

## REFERENCES

- [1] N. L. Biggs, A. T. White, "Permutation Groups and Combinatorial Structures," Cambridge Univ. Press, Cambridge, 1979.
- [2] D. Dummit, R. Foote, "Abstract Algebra," 3rd ed., J. Wiley, 2004.
- [3] T. Hungerford, "Algebra," Springer-Verlag, New York, 1980.
- [4] N. Jacobson, "Basic Algebra I," 2nd ed., W. H. Freeman, New York, 1985.
- [5] S. Lang, "Algebra," revised 3rd ed., Springer-Verlag, New York, 2002.
- [6] J. Rotman, "Advanced Modern Algebra," Prentice-Hall, Upper Saddle River, 2002.
- [7] W. R. Scott, "Group Theory," Dover, New York, 1987.
- [8] M. Suzuki, "Group Theory I," Springer-Verlag, Berlin, 1982.
- [9] R. Wilson, "The Finite Simple Groups," Springer-Verlag, New York, 2009,