

# THE CHINESE REMAINDER THEOREM

KEITH CONRAD

*We should thank the Chinese for their wonderful remainder theorem.*

Glenn Stevens

## 1. INTRODUCTION

The Chinese remainder theorem says we can uniquely solve any pair of congruences that have relatively prime moduli.

**Theorem 1.1.** *Let  $m$  and  $n$  be relatively prime positive integers. For any integers  $a$  and  $b$ , the pair of congruences*

$$x \equiv a \pmod{m}, \quad x \equiv b \pmod{n}$$

*has a solution, and this solution is uniquely determined modulo  $mn$ .*

What is important here is that  $m$  and  $n$  are relatively prime. There are *no* constraints at all on  $a$  and  $b$ .

**Example 1.2.** The congruences  $x \equiv 6 \pmod{9}$  and  $x \equiv 4 \pmod{11}$  hold when  $x = 15$ , and more generally when  $x \equiv 15 \pmod{99}$ , and they do not hold for any other  $x$ . The modulus 99 is  $9 \cdot 11$ .

We will prove the Chinese remainder theorem, including a version for more than two moduli, and see some ways it is applied to count solutions of congruences.

## 2. A PROOF OF THE CHINESE REMAINDER THEOREM

*Proof.* First we show there is always a solution. Then we will show it is unique modulo  $mn$ .

**Existence of Solution.** To show that the simultaneous congruences

$$x \equiv a \pmod{m}, \quad x \equiv b \pmod{n}$$

have a common solution in  $\mathbf{Z}$ , we give two proofs.

First proof: Write the first congruence as an equation in  $\mathbf{Z}$ , say  $x = a + my$  for some  $y \in \mathbf{Z}$ . Then the second congruence is the same as

$$a + my \equiv b \pmod{n}.$$

Subtracting  $a$  from both sides, we need to solve for  $y$  in

$$(2.1) \quad my \equiv b - a \pmod{n}.$$

Since  $(m, n) = 1$ , we know  $m \pmod{n}$  is invertible. Let  $m'$  be an inverse for  $m \pmod{n}$ , so  $mm' \equiv 1 \pmod{n}$ . Multiplying through (2.1) by  $m'$ , we have  $y \equiv m'(b - a) \pmod{n}$ , so  $y \equiv m'(b - a) + nz$  where  $z \in \mathbf{Z}$ . Then

$$x = a + my = a + m(m'(b - a) + nz) = a + mm'(b - a) + mnz.$$

So if  $x$  satisfies the original two congruences it must have this form. Let's now check this expression, for every  $z \in \mathbf{Z}$ , really satisfies the original two congruences:

$$a + mm'(b - a) + mnz \equiv a + 0 + 0 \equiv a \pmod{m}$$

and

$$a + mm'(b - a) + mnz \equiv a + 1(b - a) + 0 \equiv b \pmod{n}.$$

Second proof: Write both congruences as equations in  $\mathbf{Z}$ :  $x = a + my$  and  $x = b + nz$  for integers  $y$  and  $z$  that need to be determined. (Why would it be a bad idea to write  $x = a + my$  and  $x = b + ny$ ?) The integers of the form  $a + my$  are the numbers that are congruent to  $a \pmod{m}$ , and the integers of the form  $b + nz$  are the numbers that are congruent to  $b \pmod{n}$ . Finding a common solution to the two congruences amounts to finding  $y$  and  $z$  in  $\mathbf{Z}$  such that

$$a + my = b + nz,$$

which is the same as

$$(2.2) \quad my - nz = b - a.$$

Can we find such  $y$  and  $z$  for any  $a, b, m$ , and  $n$  where  $(m, n) = 1$ ? Bezout's identity tells us 1 is a  $\mathbf{Z}$ -linear combination of  $m$  and  $n$ , and therefore any integer is a  $\mathbf{Z}$ -linear combination of  $m$  and  $n$  (why?). Therefore integers  $y$  and  $z$  satisfying (2.2) exist.

**Uniqueness of Solution.** If  $x = c$  and  $x = c'$  both satisfy

$$x \equiv a \pmod{m}, \quad x \equiv b \pmod{n},$$

then we have  $c \equiv c' \pmod{m}$  and  $c \equiv c' \pmod{n}$ . Then  $m \mid (c - c')$  and  $n \mid (c - c')$ . Since  $(m, n) = 1$ , the product  $mn$  divides  $c - c'$ , which means  $c \equiv c' \pmod{mn}$ . This shows any two solutions to the initial pair of congruences are the same modulo  $mn$ .  $\square$

### 3. EXTENSION TO MORE THAN TWO CONGRUENCES

The Chinese remainder theorem can be extended from two congruences to any finite number of congruences, but we have to be careful about the way in which the moduli are relatively prime. Consider the three congruences

$$x \equiv 1 \pmod{6}, \quad x \equiv 4 \pmod{10}, \quad x \equiv 7 \pmod{15}.$$

While there is no common factor of 6, 10, and 15 greater than 1, these congruences do not admit a common solution: any solution to the first congruence is odd, while any solution to the second congruence is even. When we have more than two moduli, we have to be sensitive to the difference between saying numbers are collectively relatively prime (no common factor greater than 1 divides them all) and pairwise relatively prime (no common factor greater than 1 divides any two of the numbers). For instance, 6, 10, and 15 are collectively relatively prime but not pairwise relatively prime. Here is a more general form of the Chinese remainder theorem.

**Theorem 3.1.** *For  $r \geq 2$ , let  $m_1, m_2, \dots, m_r$  be nonzero integers that are pairwise relatively prime:  $(m_i, m_j) = 1$  for  $i \neq j$ . Then, for any integers  $a_1, a_2, \dots, a_r$ , the system of congruences*

$$x \equiv a_1 \pmod{m_1}, \quad x \equiv a_2 \pmod{m_2}, \quad \dots, \quad x \equiv a_r \pmod{m_r},$$

*has a solution, and this solution is uniquely determined modulo  $m_1 m_2 \cdots m_r$ .*

**Example 3.2.** The congruences  $x \equiv 1 \pmod{3}$ ,  $x \equiv 2 \pmod{5}$ ,  $x \equiv 2 \pmod{7}$  are satisfied when  $x = 37$ , more generally for any  $x \equiv 37 \pmod{105}$  and for no other  $x$ . Note  $105 = 3 \cdot 5 \cdot 7$ .

*Proof.* First we show there is always a solution. Then we will show it is unique modulo  $m_1 m_2 \cdots m_r$ .

**Existence of Solution.** We argue by induction on  $r$ . The base case  $r = 2$  is Theorem 1.1, which has been proved already.

Now we pass to the inductive step. Suppose all simultaneous congruences with  $r$  pairwise relatively prime moduli can be solved. Consider a system of simultaneous congruences with  $r + 1$  pairwise relatively prime moduli:

$$x \equiv a_1 \pmod{m_1}, \quad \dots, \quad x \equiv a_r \pmod{m_r}, \quad x \equiv a_{r+1} \pmod{m_{r+1}},$$

where  $(m_i, m_j) = 1$  for all  $i \neq j$  and the  $a_i$ 's are arbitrary. By the inductive hypothesis, there is a solution  $b$  to the first  $r$  congruences, say

$$b \equiv a_1 \pmod{m_1}, \quad b \equiv a_2 \pmod{m_2}, \quad \dots, \quad b \equiv a_r \pmod{m_r}.$$

Now consider the system of *two* congruences

$$(3.1) \quad x \equiv b \pmod{m_1 m_2 \cdots m_r}, \quad x \equiv a_{r+1} \pmod{m_{r+1}}.$$

Since  $(m_i, m_{r+1}) = 1$  for  $i = 1, 2, \dots, r$ , we have  $(m_1 m_2 \cdots m_r, m_{r+1}) = 1$ , so the two moduli in (3.1) are relatively prime. Then by the case of two congruences, namely Theorem 1.1, there is a solution to (3.1). Call it  $c$ . Since  $c \equiv b \pmod{m_1 m_2 \cdots m_r}$ , we have  $c \equiv b \pmod{m_i}$  for  $i = 1, 2, \dots, r$ . From the choice of  $b$  we have  $b \equiv a_i \pmod{m_i}$  for  $i = 1, 2, \dots, r$ . Therefore  $c \equiv a_i \pmod{m_i}$  for  $i = 1, 2, \dots, r$ . Also,  $c \equiv a_{r+1} \pmod{m_{r+1}}$  from the choice of  $c$ , so we see  $c$  satisfies the  $r + 1$  given congruences.

This concludes the inductive step, so a solution exists.

**Uniqueness of Solution.** If  $x = c$  and  $x = c'$  both satisfy

$$x \equiv a_1 \pmod{m_1}, \quad x \equiv a_2 \pmod{m_2}, \quad \dots, \quad x \equiv a_r \pmod{m_r},$$

then we have  $c \equiv c' \pmod{m_i}$  for  $i = 1, 2, \dots, r$ , so  $m_i \mid (c - c')$  for  $i = 1, 2, \dots, r$ . Since the  $m_i$ 's are pairwise relatively prime, their product  $m_1 m_2 \cdots m_r$  divides  $c - c'$ , which means  $c \equiv c' \pmod{m_1 m_2 \cdots m_r}$ . This shows any two solutions to the given system of congruences are the same when viewed modulo  $m_1 m_2 \cdots m_r$ .  $\square$

#### 4. APPLICATIONS

The significance of the Chinese remainder theorem is that it often reduces a question about modulus  $mn$ , where  $(m, n) = 1$ , to the same question for modulus  $m$  and  $n$  separately. In this way, questions about modular arithmetic can often be reduced to the special case of prime power moduli. We will see how this works for several counting problems, often using two features of modular arithmetic with two moduli:

- if  $d \mid m$  it makes sense to reduce integers mod  $m$  to integers mod  $d$ : if  $x \equiv y \pmod{m}$  then  $x \equiv y \pmod{d}$ . For example, if  $x \equiv y \pmod{10}$  then  $x \equiv y \pmod{5}$  since if  $x - y$  is divisible by 10 then it is also divisible by 5. (In contrast, it makes no sense to reduce  $x \pmod{10}$  to  $x \pmod{3}$ , since there are congruent numbers mod 10 that are incongruent mod 3, such as 5 and 15.)
- if  $x \equiv y \pmod{m}$  and  $x \equiv y \pmod{n}$  and  $(m, n) = 1$  then  $x \equiv y \pmod{mn}$ . This was used in the uniqueness part of the proof of the Chinese remainder theorem.

Our first application is to counting units.

**Theorem 4.1.** For relatively prime positive integers  $m$  and  $n$ ,  $\varphi(mn) = \varphi(m)\varphi(n)$ .

*Proof.* We work with the sets

$$U_m = \{a \bmod m, (a, m) = 1\}, \quad U_n = \{b \bmod n, (b, n) = 1\},$$

$$U_{mn} = \{c \bmod mn, (c, mn) = 1\}.$$

Then  $|U_m| = \varphi(m)$ ,  $|U_n| = \varphi(n)$ , and  $|U_{mn}| = \varphi(mn)$ . To show  $\varphi(mn) = \varphi(m)\varphi(n)$ , we will write down a bijection between  $U_{mn}$  and  $U_m \times U_n$ , which implies the two sets have the same size, and that is what the theorem is saying (since  $|U_m \times U_n| = \varphi(m)\varphi(n)$ ).

Let  $f: U_{mn} \rightarrow U_m \times U_n$  by the rule

$$f(c \bmod mn) = (c \bmod m, c \bmod n).$$

For  $c \in U_{mn}$ , we have  $(c, mn) = 1$ , so  $(c, m)$  and  $(c, n)$  equal 1, so  $c \bmod m$  and  $c \bmod n$  are units. Let's stop for a moment to take a look at an example of this function.

Take  $m = 3$  and  $n = 5$ :  $U_3 = \{1, 2\}$ ,  $U_5 = \{1, 2, 3, 4\}$ , and  $U_{15} = \{1, 2, 4, 7, 8, 11, 13, 14\}$ . The following table shows the values of the function  $f$  on each number in  $U_{15}$ . Notice that the values fill up all of  $U_3 \times U_5$  without repetition.

$c \bmod 15$	$f(c \bmod 15)$
1	(1, 1)
2	(2, 2)
4	(4, 4) = (1, 4)
7	(7, 7) = (1, 2)
8	(8, 8) = (2, 3)
11	(11, 11) = (2, 1)
13	(13, 13) = (1, 3)
14	(14, 14) = (2, 4)

There are 2 units modulo 3 and 4 units modulo 5, leading to 8 ordered pairs of units modulo 3 and units modulo 5: (1,1), (1,2), (1,3), (1,4), (2,1), (2,2), (2,3), and (2,4). All these pairs show up (and just once) in the second column of the table.

We return to the general situation and show  $f: U_{mn} \rightarrow U_m \times U_n$  is a bijection.

To see that  $f$  is one-to-one, suppose  $f(k \bmod mn) = f(\ell \bmod mn)$ . Then  $k \equiv \ell \pmod m$  and  $k \equiv \ell \pmod n$ , so since  $(m, n) = 1$  (aha!), we have  $k \equiv \ell \pmod{mn}$ . That means  $k = \ell$  in  $U_{mn}$ , so  $f$  is one-to-one.

Now we show  $f$  is onto. Pick any pair  $(a \bmod m, b \bmod n) \in U_m \times U_n$ . By the Chinese remainder theorem we can solve  $c \equiv a \pmod m$  and  $c \equiv b \pmod n$  for a  $c \in \mathbf{Z}$ . Is  $(c, mn) = 1$ ? Since  $a \bmod m$  is a unit and  $c \equiv a \pmod m$ ,  $c \bmod m$  is a unit so  $(c, m) = 1$ . Since  $b \bmod n$  is a unit and  $c \equiv b \pmod n$ ,  $c \bmod n$  is a unit so  $(c, n) = 1$ . From  $(c, m) = 1$  and  $(c, n) = 1$  we get  $(c, mn) = 1$ , so  $c \in U_{mn}$ . From the congruence conditions on  $c$ , we have  $f(c) = (a, b)$ .  $\square$

**Corollary 4.2.** *For a positive integer  $m$ ,*

$$\varphi(m) = m \prod_{p|m} \left(1 - \frac{1}{p}\right),$$

where the product runs over the primes  $p$  dividing  $m$ .

*Proof.* The formula is clear for  $m = 1$  (interpreting an empty product as 1).

Now suppose  $m > 1$ , and factor  $m$  into prime powers:

$$m = p_1^{e_1} p_2^{e_2} \cdots p_r^{e_r}.$$

The  $p_i^{e_i}$ 's are pairwise relatively prime. By an extension of Theorem 4.1 from two relatively prime terms to any number of pairwise relatively prime terms (just induct on the number of terms), we have

$$\varphi(m) = \varphi(p_1^{e_1})\varphi(p_2^{e_2}) \cdots \varphi(p_r^{e_r}).$$

Now using the formula for  $\varphi$  on prime powers,

$$\begin{aligned} \varphi(m) &= p_1^{e_1-1}(p_1-1)p_2^{e_2-1}(p_2-1) \cdots p_r^{e_r-1}(p_r-1) \\ &= p_1^{e_1} \left(1 - \frac{1}{p_1}\right) p_2^{e_2} \left(1 - \frac{1}{p_2}\right) \cdots p_r^{e_r} \left(1 - \frac{1}{p_r}\right) \\ &= m \prod_{p|m} \left(1 - \frac{1}{p}\right). \end{aligned}$$

□

**Example 4.3.** To compute  $\varphi(540) = \varphi(2^2 \cdot 3^3 \cdot 5)$ , we have

$$\begin{aligned} \varphi(540) &= 540 \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{5}\right) \\ &= 540 \cdot \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{4}{5} \\ &= 18 \cdot 8 \\ &= 144. \end{aligned}$$

An alternate calculation is

$$\begin{aligned} \varphi(540) &= \varphi(4)\varphi(27)\varphi(5) \\ &= (4-2)(27-9)(5-1) \\ &= 2 \cdot 18 \cdot 4 \\ &= 144. \end{aligned}$$

We now leave units mod  $m$  and look at squares mod  $m$ .

**Theorem 4.4.** For  $m \in \mathbf{Z}^+$  with  $m \geq 2$ , let  $S_m = \{x^2 \bmod m\}$  be the set of squares modulo  $m$ . When  $(m, n) = 1$ ,  $|S_{mn}| = |S_m| \cdot |S_n|$ .

Note  $S_m$  is *all* squares modulo  $m$ , including 0. So  $S_5 = \{0, 1, 4\}$ , for example.

*Proof.* We will use the Chinese remainder theorem *twice*.

If  $a \equiv x^2 \bmod mn$  then  $a \equiv x^2 \bmod m$  and  $a \equiv x^2 \bmod n$ . Thus any square modulo  $mn$  reduces to a square modulo  $m$  and a square modulo  $n$ . So we have a function  $f: S_{mn} \rightarrow S_m \times S_n$  by  $f(a \bmod mn) = (a \bmod m, a \bmod n)$ . Let's take a look at an example.

Set  $m = 3$  and  $n = 5$ , so  $S_3 = \{0, 1\}$ ,  $S_5 = \{0, 1, 4\}$  and  $S_{15} = \{0, 1, 4, 6, 9, 10\}$ . The table below gives the values of  $f$  on  $S_{15}$ . The values fill up  $S_3 \times S_5$  without repetition.

$c \bmod 15$	$f(c \bmod 15)$
0	(0, 0)
1	(1, 1)
4	(4, 4) = (1, 4)
6	(6, 6) = (0, 1)
9	(9, 9) = (0, 4)
10	(10, 10) = (1, 0)

Returning to the general case, to show  $f$  is one-to-one let's suppose  $f(c \bmod mn) = f(c' \bmod mn)$ . Then  $c \equiv c' \pmod m$  and  $c \equiv c' \pmod n$ , so  $c \equiv c' \pmod mn$  since  $(m, n) = 1$ .

To show  $f$  is onto, pick a pair of squares  $b \bmod m$  and  $c \bmod n$ , say  $b \equiv y^2 \pmod m$  and  $c \equiv z^2 \pmod n$ . By the Chinese remainder theorem, there is an integer  $a$  satisfying

$$a \equiv b \pmod m, \quad a \equiv c \pmod n.$$

We want to say  $f(a) = (b, c)$ , but is  $a \bmod mn$  a square? From the expressions for  $b \bmod m$  and  $c \bmod n$  as squares,  $a \equiv y^2 \pmod m$  and  $a \equiv z^2 \pmod n$ , but  $y$  and  $z$  are not related to each other. They certainly don't have to be the same integer, so these two congruences on their own don't tell us  $a \bmod mn$  is a square. Using the Chinese remainder theorem *again*, however, there is  $x \in \mathbf{Z}$  such that

$$x \equiv y \pmod m, \quad x \equiv z \pmod n,$$

so  $x^2 \equiv y^2 \pmod m$  and  $x^2 \equiv z^2 \pmod n$ . Therefore  $a \equiv x^2 \pmod m$  and  $a \equiv x^2 \pmod n$ , so  $a \equiv x^2 \pmod mn$ , so  $a \bmod mn$  is in fact a square. Thus  $a \in S_{mn}$  and  $f(a) = (b, c)$ .  $\square$

**Example 4.5.** For a prime  $p$ , the number of nonzero squares mod  $p$  is  $(p-1)/2$  and 0 is a square, so the total number of squares mod  $p$  is  $1 + (p-1)/2 = (p+1)/2$ . Thus  $|S_p| = (p+1)/2$ . So if  $n = p_1 p_2 \dots p_r$  is squarefree,  $|S_n| = |S_{p_1}| \dots |S_{p_r}| = \frac{p_1+1}{2} \dots \frac{p_r+1}{2}$ . If  $n = p_1^{e_1} \dots p_r^{e_r}$ , we have  $|S_n| = |S_{p_1^{e_1}}| \dots |S_{p_r^{e_r}}|$ , so a formula for  $|S_{p^e}|$  when  $e > 1$  (which we don't give here) would lead to a formula for  $|S_m|$  in general.

We turn now from counting all the squares mod  $m$  to counting how often something is a square mod  $m$ .

**Example 4.6.** We can write  $1 \bmod 15$  as a square in *four* ways:  $1 \equiv 1^2 \equiv 4^2 \equiv 9^2 \equiv 14^2 \pmod{15}$ .

**Theorem 4.7.** Let  $m \in \mathbf{Z}^+$  have prime factorization  $p_1^{e_1} \dots p_r^{e_r}$ . For any integer  $a$ , the congruence  $x^2 \equiv a \pmod m$  is solvable if and only if the separate congruences  $x^2 \equiv a \pmod{p_i^{e_i}}$  are solvable for  $i = 1, 2, \dots, r$ .

Furthermore, if the congruence  $x^2 \equiv a \pmod{p_i^{e_i}}$  has  $N_i$  solutions, then the congruence  $x^2 \equiv a \pmod m$  has  $N_1 N_2 \dots N_r$  solutions.

**Example 4.8.** The congruences  $x^2 \equiv 1 \pmod 3$  and  $x^2 \equiv 1 \pmod 5$  each have two solutions, so  $x^2 \equiv 1 \pmod{15}$  has  $2 \cdot 2 = 4$  solutions; we saw the four square roots of 1 mod 15 before the statement of Theorem 4.7.

*Proof.* If  $x \in \mathbf{Z}$  satisfies  $x^2 \equiv a \pmod m$ , then  $x^2 \equiv a \pmod{p_i^{e_i}}$  for all  $i$ .

Conversely, suppose each of the congruences  $x^2 \equiv a \pmod{p_i^{e_i}}$  has a solution, say  $x_i^2 \equiv a \pmod{p_i^{e_i}}$  for some integers  $x_i$ . Since the  $p_i^{e_i}$ 's are pairwise relatively prime, the Chinese remainder theorem tells us there is an  $x$  such that  $x \equiv x_i \pmod{p_i^{e_i}}$  for all  $i$ . Then  $x^2 \equiv x_i^2 \pmod{p_i^{e_i}}$  for all  $i$ , so  $x^2 \equiv a \pmod{p_i^{e_i}}$  for all  $i$ . Since  $x^2 - a$  is divisible by each  $p_i^{e_i}$  it is divisible by  $m$ , so  $x^2 \equiv a \pmod m$ .

To count the solutions modulo  $m$ , we again use the Chinese remainder theorem. Any choice of solution  $x_i \pmod{p_i^{e_i}}$  for each  $i$  fits together in exactly one way to a number  $x \pmod m$ , and this number will satisfy  $x^2 \equiv a \pmod m$ . Therefore we can count solutions modulo  $m$  by counting solutions modulo each  $p_i^{e_i}$  and multiply the counts thanks to the independence of the choice of solutions for different primes.  $\square$

**Example 4.9.** To decide if 61 is a square modulo 75, we check whether 61 is a square modulo 3 and modulo 25. Since  $61 \equiv 1 \pmod{3}$ , it is a square modulo 3. Since  $61 \equiv 11 \equiv 6^2 \pmod{25}$ , it is a square modulo 25. Therefore 61 is a square modulo 75. In fact, we can get a square root by solving the congruences

$$x \equiv 1 \pmod{3}, \quad x \equiv 6 \pmod{25}.$$

A solution is  $x = 31$ , so  $61 \equiv 31^2 \pmod{75}$ .

If you scrutinize the proofs of Theorems 4.4 and 4.7 to see why it was important we were working with squares, you'll see that it really wasn't; the only thing that really matters is that squaring is a polynomial expression. With this in mind, we get the following generalizations from squares to values of other polynomials.

**Theorem 4.10.** *Let  $f(x)$  be any polynomial with integer coefficients. For a positive integer  $m \geq 2$ , let  $N_f(m) = |\{f(x) \pmod{m} : 0 \leq x \leq m-1\}|$  be the number of values of  $f$  on different integers mod  $m$ . If  $m$  has prime factorization*

$$m = p_1^{e_1} \cdots p_r^{e_r},$$

*we have  $N_f(m) = N_f(p_1^{e_1}) \cdots N_f(p_r^{e_r})$ .*

*Proof.* Proceed as in the proof of Theorem 4.4, which is the special case  $f(x) = x^2$ .  $\square$

**Theorem 4.11.** *Let  $f(x)$  be any polynomial with integer coefficients. For a positive integer  $m$  with prime factorization*

$$m = p_1^{e_1} \cdots p_r^{e_r},$$

*the congruence  $f(x) \equiv 0 \pmod{m}$  is solvable if and only if the congruences  $f(x) \equiv 0 \pmod{p_i^{e_i}}$  are each solvable.*

*Moreover, if  $f(x) \equiv 0 \pmod{p_i^{e_i}}$  has  $N_i$  solutions, then the congruence  $f(x) \equiv 0 \pmod{m}$  has  $N_1 N_2 \cdots N_r$  solutions.*

*Proof.* Argue as in the proof of Theorem 4.7, which is the special case  $f(x) = x^2 - a$ .  $\square$

Theorem 4.11 tells us that finding solutions to a polynomial equation modulo positive integers is reduced by the Chinese remainder theorem to the case of understanding solutions modulo prime powers.

Consider now the following situation:  $f(x)$  is a polynomial with integral coefficients and every value  $f(n)$ , for  $n \in \mathbf{Z}$ , is either a multiple of 2 or a multiple of 3. For instance, if  $f(x) = x^2 - x$  then  $f(n) = n^2 - n$  is even for all  $n$ . Or if  $f(x) = x^3 - x$  then  $f(n) = n^3 - n$  is a multiple of 3 for all  $n$ . But these examples are kind of weak: what about a mixed example where every  $f(n)$  is a multiple of 2 or 3 but some  $f(n)$  are multiples of 2 and not 3 while other  $f(n)$  are multiples of 3 and not 2? Actually, no such polynomial exists! The only way  $f(n)$  can be divisible either by 2 or 3 for all  $n$  is if it is a multiple of 2 for all  $n$  or a multiple of 3 for all  $n$ . To explain this, we will use the Chinese remainder theorem.

**Theorem 4.12.** *Let  $f(x)$  be a polynomial with integral coefficients. Suppose there is a finite set of primes  $p_1, \dots, p_r$  such that, for every integer  $n$ ,  $f(n)$  is divisible by some  $p_i$ . Then there is one  $p_i$  such that  $p_i \mid f(n)$  for every  $n \in \mathbf{Z}$ .*

*Proof.* Suppose the conclusion is false. Then, for each  $p_i$ , there is an  $a_i \in \mathbf{Z}$  such that  $p_i$  does not divide  $f(a_i)$ . Said differently,  $f(a_i) \not\equiv 0 \pmod{p_i}$ .

Since the  $p_i$ 's are different primes, we can use the Chinese remainder theorem to find a single integer  $a$  such that  $a \equiv a_i \pmod{p_i}$  for  $i = 1, 2, \dots, r$ . Then  $f(a) \equiv f(a_i) \pmod{p_i}$  for

$i = 1, 2, \dots, r$  (why?), so  $f(a) \not\equiv 0 \pmod{p_i}$  for all  $i$ . However, the assumption in the theorem was that every value of the polynomial on integers is divisible by some  $p_i$ , so we have a contradiction.  $\square$

**Remark 4.13.** It is natural to believe an analogous result for divisibility by squares of primes. Specifically, if  $f(x)$  is a polynomial with integral coefficients and there is a finite set of primes  $p_1, \dots, p_r$  such that, for every integer  $n$ ,  $f(n)$  is divisible by some  $p_i^2$ , then there should be one  $p_i$  such that  $p_i^2 \mid f(n)$  for every  $n \in \mathbf{Z}$ . If you try to adapt the proof of Theorem 4.12 to this setting, it breaks down (where?). While this analogue for divisibility by squares of primes is plausible, it is still an open problem as far as I am aware.

Our final application of the Chinese remainder theorem is to an interpolation problem. Given  $n$  points in the plane,  $(a_1, b_1), \dots, (a_n, b_n)$ , with the  $a_i$ 's distinct, we would like to find a polynomial  $f(T)$  in  $\mathbf{R}[T]$  whose graph passes through these points:  $f(a_i) = b_i$  for  $i = 1, 2, \dots, n$ . This task can be converted to a set of simultaneous congruences in  $\mathbf{R}[T]$ , which can be solved using the Chinese remainder theorem in  $\mathbf{R}[T]$ , not  $\mathbf{Z}$ . First let's state the Chinese remainder theorem for polynomials.

**Theorem 4.14.** *For  $r \geq 2$ , let  $m_1(T), m_2(T), \dots, m_r(T)$  be nonzero polynomials in  $\mathbf{R}[T]$  which are pairwise relatively prime:  $(m_i(T), m_j(T)) = 1$  for  $i \neq j$ . Then, for any polynomials  $a_1(T), a_2(T), \dots, a_r(T)$ , the system of congruences*

$$f(T) \equiv a_1(T) \pmod{m_1(T)}, \quad f(T) \equiv a_2(T) \pmod{m_2(T)}, \quad \dots, \quad f(T) \equiv a_r(T) \pmod{m_r(T)},$$

*has a solution  $f(T)$  in  $\mathbf{R}[T]$ , and this solution is unique modulo  $m_1(T)m_2(T)\cdots m_r(T)$ .*

The proof of this is identical to that of the Chinese remainder theorem for  $\mathbf{Z}$ , so we leave it to the reader as an exercise.

**Theorem 4.15.** *In  $\mathbf{R}$ , pick  $n$  distinct numbers  $a_1, a_2, \dots, a_n$  and any numbers  $b_1, b_2, \dots, b_n$ . There is a unique polynomial  $f(T)$  of degree  $< n$  in  $\mathbf{R}[T]$ , possibly 0, such that  $f(a_i) = b_i$  for all  $i$ .*

*Proof.* To say  $f(a_i) = b_i$  is the same as  $f(T) \equiv b_i \pmod{T - a_i}$  (why?). Consider the system of congruences

$$f(T) \equiv b_1 \pmod{T - a_1}, \quad f(T) \equiv b_2 \pmod{T - a_2}, \quad \dots, \quad f(T) \equiv b_n \pmod{T - a_n}$$

for an unknown  $f(T)$  in  $\mathbf{R}[T]$ . Since the  $a_i$ 's are *distinct*, the polynomials  $T - a_1, \dots, T - a_n$  are pairwise relatively prime in  $\mathbf{R}[T]$ . Therefore, by the Chinese remainder theorem in  $\mathbf{R}[T]$ , there is an  $f(T)$  in  $\mathbf{R}[T]$  satisfying all of the above congruences. It follows that  $f(a_i) = b_i$  for all  $i$ .

We have no initial control over  $\deg f$  for the common solution  $f$ . However, since we can adjust  $f(T)$  modulo  $(T - a_1)\cdots(T - a_n)$  without changing the congruence conditions, we can replace  $f(T)$  with its remainder under division by  $(T - a_1)\cdots(T - a_n)$ , which has degree  $n$ . Then  $\deg f < n$  with  $f(a_i) = b_i$  for all  $i$ .

We have shown a desired  $f(T)$  exists. To see it is unique, suppose  $f_1(T)$  and  $f_2(T)$  both have degree less than  $n$  and satisfy

$$f(T) \equiv b_1 \pmod{T - a_1}, \quad f(T) \equiv b_2 \pmod{T - a_2}, \quad \dots, \quad f(T) \equiv b_n \pmod{T - a_n}.$$

Then, by the uniqueness in the Chinese remainder theorem, we have

$$f_1(T) \equiv f_2(T) \pmod{(T - a_1)\cdots(T - a_n)}.$$

Since  $f_1(T)$  and  $f_2(T)$  have degree less than  $n$ , this congruence modulo a polynomial of degree  $n$  implies  $f_1(T) = f_2(T)$  in  $\mathbf{R}[T]$ .  $\square$

The fact that polynomial interpolation is identical to solving a system of polynomial congruences (with linear moduli) suggests that we should think about solving a system of integer congruences as *arithmetic* interpolation.

There is nothing essential about  $\mathbf{R}$  in Theorem 4.15 except that it's a field. The Chinese remainder theorem goes through for  $F[T]$  with  $F$  any field, not just  $\mathbf{R}$ , and Theorem 4.15 carries over to any field:

**Theorem 4.16.** *Let  $F$  be any field. For  $n$  distinct numbers  $a_1, a_2, \dots, a_n$  in  $F$  and any numbers  $b_1, b_2, \dots, b_n$  in  $F$ , there is a unique polynomial  $f(T)$  of degree  $< n$  in  $F[T]$ , possibly 0, such that  $f(a_i) = b_i$  for all  $i$ .*

The proof is identical to that of Theorem 4.15.